



US 20030228594A1

(19) **United States**

(12) **Patent Application Publication**
Koster

(10) **Pub. No.: US 2003/0228594 A1**

(43) **Pub. Date: Dec. 11, 2003**

(54) **DNA DIAGNOSTICS BASED ON MASS SPECTROMETRY**

(76) Inventor: **Hubert Koster**, Lugano-Cassarate (CH)

Correspondence Address:

Stephanie Seidman

Heller Ehrman White & McAuliffe LLP

7th Floor

4350 La Jolla Village Drive

San Diego, CA 92122-1246 (US)

(21) Appl. No.: **10/375,714**

(22) Filed: **Feb. 24, 2003**

Related U.S. Application Data

(63) Continuation of application No. 09/879,341, filed on Jun. 11, 2001, now Pat. No. 6,589,485.

Continuation of application No. 09/796,416, filed on Feb. 28, 2001, now Pat. No. 6,500,621, which is a continuation of application No. 09/495,444, filed on

Jan. 31, 2000, now Pat. No. 6,300,076, and which is a continuation of application No. 09/504,245, filed on Feb. 15, 2000, now Pat. No. 6,221,605.

Continuation of application No. 09/287,679, filed on Apr. 6, 1999, now Pat. No. 6,258,538.

Continuation of application No. 08/617,256, filed on Mar. 18, 1996, now Pat. No. 6,043,031.

Continuation-in-part of application No. 08/406,199, filed on Mar. 17, 1995, now Pat. No. 5,605,798.

Said application No. 08/617,256 is a continuation-in-part of application No. 08/406,199, filed on Mar. 17, 1995, now Pat. No. 5,605,798.

Publication Classification

(51) **Int. Cl.⁷ C12Q 1/68**

(52) **U.S. Cl. 435/6**

(57)

ABSTRACT

Fast and highly accurate mass spectrometry-based processes for detecting particular nucleic acid molecules and mutations in the molecules are provided.




FIG. 1A




FIG. 1B




FIG. 1C




FIG. 2




FIG. 3





FIG. 5






FIG. 6A




FIG. 6B




FIG. 6C





FIG. 7B




FIG. 8




FIG. 9




FIG. 10A-1




FIG. 10A-3




FIG. 10A-2




FIG. 10A-4




FIG. 10B




FIG. 12A




FIG. 10C




FIG. 11




FIG. 12B




FIG. 12C




FIG. 19




FIG. 13




FIG. 14




FIG. 15




FIG. 16




FIG. 17




FIG. 18




FIG. 20A




FIG. 20B




FIG. 21A

MetaPhor Agarose Gel
3, 5 %




FIG. 21B

Polyacrylamidgel
12 %




FIG. 21C

MOLECULAR WEIGHT OF THE VARIABLE FRAGMENTS IN D₀:

			$\epsilon 2/\epsilon 2$	$\epsilon 3/\epsilon 3$	$\epsilon 4/\epsilon 4$	$\epsilon 2/\epsilon 3$	$\epsilon 2/\epsilon 4$	$\epsilon 3/\epsilon 4$
91 bp	SENSE	28421	X	X		X	X	X
	ANTISENSE	27864						
83 bp	SENSE	25747	X			X	X	
	ANTISENSE	25591						
72 bp	SENSE	22440			X		X	X
	ANTISENSE	21494						
48 bp	SENSE	14844		X	X	X	X	X
	ANTISENSE	14857						
35 bp	SENSE	10921		X	X	X	X	X
	ANTISENSE	10751						

FIG. 22A




FIG. 22B




FIG. 23A





FIG. 23B



M 1 2 3 4 5 6

FIG. 24




FIG. 25A




FIG. 25B




FIG. 25C




FIG. 26




FIG. 27




FIG. 28




FIG. 29




FIG. 30A




FIG. 30B




FIG. 31A




FIG. 31B




FIG. 32




FIG. 33A





FIG. 33B

506507508
IleIlePhe
ACCATTAAAGAAATATCATCTTTGGTGTTTCTATGATGAATATAGAACGTCATC (SEQ. ID. NO. 60)
primer ACCACAAAGGATACTACTTTATATC (7289,8) (SEQ. ID. NO. 61)
wildtype TAGAAACCCACAAAGGATACTACTTTATATC (8846,8) (SEQ. ID. NO. 62)
ΔF508 TA---ACCACAAAGGATACTACTTTATATC (7891,2) (SEQ. ID. NO. 63)
ΔI507 TAG---AAACCACAAAGGATACTACTTTATATC (8846,8) (SEQ. ID. NO. 64)

FIG. 34A

506507508
IleIlePhe
ACCATTAAAGAAATATCATCTTTGGTGTTTCTATGATGAATATAGAACGTCATC (SEQ. ID. NO. 60)
primer ACCACAAAGGATACTACTTTATATC (7289,8) (SEQ. ID. NO. 61)
wildtype CTTTATAGTAGAAACCCACAAAGGATACTACTTTATATC (11612,6) (SEQ. ID. NO. 65)
ΔF508 CTTTATAGTA---ACCACAAAGGATACTACTTTATATC (10657,0) (SEQ. ID. NO. 66)
ΔI507 CTTTATAG---AAACCACAAAGGATACTACTTTATATC (10666,0) (SEQ. ID. NO. 67)
506Ser CGTAGAAACCCACAAAGGATACTACTTTATATC (9465,2) (SEQ. ID. NO. 68)


FIG. 34B



$\Delta F508$ (7916.9/7891.2)

Homozygous

FIG. 35A



$\Delta F508$ (7913.0/7891.2)

Homozygous

FIG. 35B




FIG. 38

1 2 3 4 5 6




FIG. 39




FIG. 40A






FIG. 40B






FIG. 42A




FIG. 42B




FIG. 43A




FIG. 43B




FIG. 44A




FIG. 44B




FIG. 44C




FIG. 44D





FIG. 45






FIG. 47A




FIG. 47B




FIG. 47C




FIG. 47D

FIG. 48A




FIG. 48B




FIG. 48C





FIG. 48D



DNA DIAGNOSTICS BASED ON MASS SPECTROMETRY

RELATED APPLICATIONS

[0001] This application is a continuation of allowed U.S. application Ser. No. 09/879,341, filed Jun. 11, 2001. This application also is a continuation of U.S. application Ser. No. 09/796,416, filed Feb. 28, 2001, now U.S. Pat. No. 6,500,621, and is a continuation of U.S. application Ser. No. 09/495,444, filed Jan. 31, 2000, now U.S. Pat. No. 6,300,076, and of U.S. application Ser. No. 09/504,245, filed Feb. 15, 2000, now U.S. Pat. No. 6,221,605. This application also is a continuation of U.S. application Ser. No. 09/287,679, filed Apr. 6, 1999, now U.S. Pat. No. 6,258,538. This application is also a continuation of U.S. application Ser. No. 08/617,256, filed Mar. 18, 1996 and now U.S. Pat. No. 6,043,031. This application is also a continuation-in-part of U.S. application Ser. No. 08/406,199 filed Mar. 17, 1995, and now U.S. Pat. No. 5,605,798. U.S. application Ser. No. 08/617,256 is a continuation-in-part of U.S. application Ser. No. 08/406,199. The subject matter of each of these applications is herein incorporated by reference in its entirety.

BACKGROUND OF THE INVENTION

[0002] The genetic information of all living organisms (e.g. animals, plants and microorganisms) is encoded in deoxyribonucleic acid (DNA). In humans, the complete genome is comprised of about 100,000 genes located on 24 chromosomes (The Human Genome, T. Strachan, BIOS Scientific Publishers, 1992). Each gene codes for a specific protein which after its expression via transcription and translation, fulfills a specific biochemical function within a living cell. Changes in a DNA sequence are known as mutations and can result in proteins with altered or in some cases even lost biochemical activities; this in turn can cause genetic disease. Mutations include nucleotide deletions, insertions or alterations (i.e. point mutations). Point mutations can be either "missense", resulting in a change in the amino acid sequence of a protein or "nonsense" coding for a stop codon and thereby leading to a truncated protein.

[0003] More than 3000 genetic diseases are currently known (Human Genome Mutations, D. N. Cooper and M. Krawczak, BIOS Publishers, 1993), including hemophilias, thalassemias, Duchenne Muscular Dystrophy (DMD), Huntington's Disease (HD), Alzheimer's Disease and Cystic Fibrosis (CF). In addition to mutated genes, which result in genetic disease, certain birth defects are the result of chromosomal abnormalities such as Trisomy 21 (Down's Syndrome), Trisomy 13 (Patau Syndrome), Trisomy 18 (Edward's Syndrome), Monosomy X (Turner's Syndrome) and other sex chromosome aneuploidies such as Klinefelter's Syndrome (XXY). Further, there is growing evidence that certain DNA sequences may predispose an individual to any of a number of diseases such as diabetes, arteriosclerosis, obesity, various autoimmune diseases and cancer (e.g. colorectal, breast, ovarian, lung).

[0004] Viruses, bacteria, fungi and other infectious organisms contain distinct nucleic acid sequences, which are different from the sequence contained in the host cell. Therefore, infectious organisms can also be detected and identified based on their specific DNA sequences.

[0005] Since the sequence of about 16 nucleotides is specific on statistical grounds even for the size of the human

genome, relatively short nucleic acid sequences can be used to detect normal and defective genes in higher organisms and to detect infectious microorganisms (e.g. bacteria, fungi, protists and yeast) and viruses. DNA sequences can even serve as a fingerprint for detection of different individuals within the same species. (Thompson, J. S. and M. W. Thompson, eds., *Genetics in Medicine*, W. B. Saunders Co., Philadelphia, Pa. (1986).

[0006] Several methods for detecting DNA are currently being used. For example, nucleic acid sequences can be identified by comparing the mobility of an amplified nucleic acid fragment with a known standard by gel electrophoresis, or by hybridization with a probe, which is complementary to the sequence to be identified. Identification, however, can only be accomplished if the nucleic acid fragment is labeled with a sensitive reporter function (e.g. radioactive (^{32}P , ^{35}S), fluorescent or chemiluminescent). However, radioactive labels can be hazardous and the signals they produce decay over time. Non-isotopic labels (e.g. fluorescent) suffer from a lack of sensitivity and fading of the signal when high intensity lasers are being used. Additionally, performing labeling, electrophoresis and subsequent detection are laborious, time-consuming and error-prone procedures. Electrophoresis is particularly error-prone, since the size or the molecular weight of the nucleic acid cannot be directly correlated to the mobility in the gel matrix. It is known that sequence specific effects, secondary structures and interactions with the gel matrix are causing artifacts.

[0007] In general, mass spectrometry provides a means of "weighing" individual molecules by ionizing the molecules in vacuo and making them "fly" by volatilization. Under the influence of combinations of electric and magnetic fields, the ions follow trajectories depending on their individual mass (m) and charge (z). In the range of molecules with low molecular weight, mass spectrometry has long been part of the routine physical-organic repertoire for analysis and characterization of organic molecules by the determination of the mass of the parent molecular ion. In addition, by arranging collisions of this parent molecular ion with other particles (e.g. argon atoms), the molecular ion is fragmented forming secondary ions by the so-called collision induced dissociation (CID). The fragmentation pattern/pathway very often allows the derivation of detailed structural information. Many applications of mass spectrometric methods are known in the art, particularly in biosciences, and can be found summarized in *Methods of Enzymology*, Vol. 193: "Mass Spectrometry" (J. A. McCloskey, editor), 1990, Academic Press, New York.

[0008] Due to the apparent analytical advantages of mass spectrometry in providing high detection sensitivity, accuracy of mass measurements, detailed structural information by CID in conjunction with an MS/MS configuration and speed, as well as on-line data transfer to a computer, there has been considerable interest in the use of mass spectrometry for the structural analysis of nucleic acids. Recent reviews summarizing this field include K. H. Schram, "Mass Spectrometry of Nucleic Acid Components, Biomedical Applications of Mass Spectrometry" 34, 203-287 (1990); and P. F. Crain, "Mass Spectrometric Techniques in Nucleic Acid Research, "Mass Spectrometry Reviews 9, 505-554 (1990).

[0009] However, nucleic acids are very polar biopolymers that are very difficult to volatilize. Consequently, mass

spectrometric detection has been limited to low molecular weight synthetic oligonucleotides by determining the mass of the parent molecular ion and through this, confirming the already known oligonucleotide sequence, or alternatively, confirming the known sequence through the generation of secondary ions (fragment ions) via CID in the MS/MS configuration utilizing, in particular, for the ionization and volatilization, the method of fast atomic bombardment (FAB mass spectrometry) or plasma desorption (PD mass spectrometry). As an example, the application of FAB to the analysis of protected dimeric blocks for chemical synthesis of oligodeoxynucleotides has been described (Wolter et al. *Biomedical Environmental Mass Spectrometry* 14: 111-116 (1987)).

[0010] Two more recent ionizations/desorption techniques are electrospray/ion spray (ES) and matrix-assisted laser desorption/ionization (MALDI). ES mass spectrometry has been introduced by Yamashita et al. (*J. Phys. Chem.* 88: 4451-59 (1984); PCT Application No. WO 90/14148) and current applications are summarized in recent review articles (R. D. Smith et al., *Anal. Chem.* 62: 882-89 (1990) and B. Ardrey, *Electrospray Mass Spectrometry, Spectroscopy Europe* 4: 10-18 (1992)). The molecular weights of a tetradecanucleotide (Covey et al. "The Determination of Protein, Oligonucleotide and Peptide Molecular Weights by Ion Spray Mass Spectrometry," *Rapid Communications in Mass Spectrometry* 2: 249-256 (1988)), and of a 21-mer (*Methods in Enzymology* 193, "Mass Spectrometry" (McCloskey, editor), p. 425, 1990, Academic Press, New York) have been published. As a mass analyzer, a quadrupole is most frequently used. The determination of molecular weights in femtomole amounts of sample is very accurate due to the presence of multiple ion peaks which all could be used for the mass calculation.

[0011] MALDI mass spectrometry, in contrast, can be particularly attractive when a time-of-flight (TOF) configuration is used as a mass analyzer. The MALDI-TOF mass spectrometry has been introduced by Hillenkamp et al. ("Matrix Assisted UV-Laser Desorption/Ionization: A New Approach to Mass Spectrometry of Large Biomolecules," *Biological Mass Spectrometry* (Burlingame and McCloskey, editors), Elsevier Science Publishers, Amsterdam, pp. 49-60, 1990). Since, in most cases, no multiple molecular ion peaks are produced with this technique, the mass spectra, in principle, look simpler compared to ES mass spectrometry.

[0012] Although DNA molecules up to a molecular weight of 410,000 daltons have been desorbed and volatilized (Nelson et al., "Volatilization of High Molecular Weight DNA by Pulsed Laser Ablation of Frozen Aqueous Solutions," *Science* 246: 1585-87 (1989)), this technique has so far only shown very low resolution (oligonucleotides up to 18 nucleotides, Huth-Fehre et al., *Rapid Communications in Mass Spectrometry* 6: 209-13 (1992); DNA fragments up to 500 nucleotides in length, Tang, K. et al., *Rapid Communications in Mass Spectrometry* 8: 727-730 (1994); and a double-stranded DNA of 28 base pairs (Williams et al., "Time-of Flight Mass Spectrometry of Nucleic Acids by Laser Ablation and Ionization from a Frozen Aqueous Matrix," *Rapid Communications in Mass Spectrometry* 4: 348-351 (1990)).

[0013] Japanese Patent No. 59-131909 describes an instrument, which detects nucleic acid fragments separated either

by electrophoresis, liquid chromatography or high speed gel filtration. Mass spectrometric detection is achieved by incorporating into the nucleic acids, atoms which normally do not occur in DNA such as S, Br, I or Ag, Au, Pt, Os, Hg.

SUMMARY OF THE INVENTION

[0014] The instant invention provides mass spectrometric processes for detecting a particular nucleic acid sequence in a biological sample. Depending on the sequence to be detected, the processes can be used, for example, to diagnose (e.g. prenatally or postnatally) a genetic disease or chromosomal abnormality; a predisposition to a disease or condition (e.g. obesity, atherosclerosis, cancer), or infection by a pathogenic organism (e.g. virus, bacteria, parasite or fungus); or to provide information relating to identity, heredity, or compatibility (e.g. HLA phenotyping).

[0015] In a first embodiment, a nucleic acid molecule containing the nucleic acid sequence to be detected (i.e. the target) is initially immobilized to a solid support. Immobilization can be accomplished, for example, based on hybridization between a portion of the target nucleic acid molecule, which is distinct from the target detection site and a capture nucleic acid molecule, which has been previously immobilized to a solid support. Alternatively, immobilization can be accomplished by direct bonding of the target nucleic acid molecule and the solid support. Preferably, there is a spacer (e.g. a nucleic acid molecule) between the target nucleic acid molecule and the support. A detector nucleic acid molecule (e.g. an oligonucleotide or oligonucleotide mimetic), which is complementary to the target detection site can then be contacted with the target detection site and formation of a duplex, indicating the presence of the target detection site can be detected by mass spectrometry. In preferred embodiments, the target detection site is amplified prior to detection and the nucleic acid molecules are conditioned. In a further preferred embodiment, the target detection sequences are arranged in a format that allows multiple simultaneous detections (multiplexing), as well as parallel processing using oligonucleotide arrays ("DNA chips").

[0016] In a second embodiment, immobilization of the target nucleic acid molecule is an optional rather than a required step. Instead, once a nucleic acid molecule has been obtained from a biological sample, the target detection sequence is amplified and directly detected by mass spectrometry. In preferred embodiments, the target detection site and/or the detector oligonucleotides are conditioned prior to mass spectrometric detection. In another preferred embodiment, the amplified target detection sites are arranged in a format that allows multiple simultaneous detections (multiplexing), as well as parallel processing using oligonucleotide arrays ("DNA chips").

[0017] In a third embodiment, nucleic acid molecules which have been replicated from a nucleic acid molecule obtained from a biological sample can be specifically digested using one or more nucleases (using deoxyribonucleases for DNA or ribonucleases for RNA) and the fragments captured on a solid support carrying the corresponding complementary sequences. Hybridization events and the actual molecular weights of the captured target sequences provide information on whether and where mutations in the gene are present. The array can be analyzed spot by spot using mass spectrometry. DNA can be similarly digested

using a cocktail of nucleases including restriction endonucleases. In a preferred embodiment, the nucleic acid fragments are conditioned prior to mass spectrometric detection.

[0018] In a fourth embodiment, at least one primer with 3' terminal base complementarity to an allele (mutant or normal) is hybridized with a target nucleic acid molecule, which contains the allele. An appropriate polymerase and a complete set of nucleoside triphosphates or only one of the nucleoside triphosphates are used in separate reactions to furnish a distinct extension of the primer. Only if the primer is appropriately annealed (i.e. no 3' mismatch) and if the correct (i.e. complementary) nucleotide is added, will the primer be extended. Products can be resolved by molecular weight shifts as determined by mass spectrometry.

[0019] In a fifth embodiment, a nucleic acid molecule containing the nucleic acid sequence to be detected (i.e. the target) is initially immobilized to a solid support. Immobilization can be accomplished, for example, based on hybridization between a portion of the target nucleic acid molecule, which is distinct from the target detection site and a capture nucleic acid molecule, which has been previously immobilized to a solid support. Alternatively, immobilization can be accomplished by direct bonding of the target nucleic acid molecule and the solid support. Preferably, there is a spacer (e.g. a nucleic acid molecule) between the target nucleic acid molecule and the support. A nucleic acid molecule that is complementary to a portion of the target detection site that is immediately 5' of the site of a mutation is then hybridized with the target nucleic acid molecule. The addition of a complete set of dideoxynucleosides or 3'-dideoxynucleoside triphosphates (e.g. pppAdd, pppTdd, pppCdd and pppGdd) and a DNA dependent DNA polymerase allows for the addition only of the one dideoxynucleoside or 3'-dideoxynucleoside triphosphate that is complementary to X. The hybridization product can then be detected by mass spectrometry.

[0020] In a sixth embodiment, a target nucleic acid is hybridized with a complementary oligonucleotides that hybridize to the target within a region that includes a mutation M. The heteroduplex is then contacted with an agent that can specifically cleave at an unhybridized portion (e.g. a single strand specific endonuclease), so that a mismatch, indicating the presence of a mutation, results in a cleavage of the target nucleic acid. The two cleavage products can then be detected by mass spectrometry.

[0021] In a seventh embodiment, which is based on the ligase chain reaction (LCR), a target nucleic acid is hybridized with a set of ligation educts and a thermostable DNA ligase, so that the ligase educts become covalently linked to each other, forming a ligation product. The ligation product can then be detected by mass spectrometry and compared to a known value. If the reaction is performed in a cyclic manner, the ligation product obtained can be amplified to better facilitate detection of small volumes of the target nucleic acid. Selection between wildtype and mutated primers at the ligation point can result in a detection of a point mutation.

[0022] The processes of the invention provide for increased accuracy and reliability of nucleic acid detection by mass spectrometry. In addition, the processes allow for rigorous controls to prevent false negative or positive results. The processes of the invention avoid electrophoretic

steps, labeling and subsequent detection of a label. In fact it is estimated that the entire procedure, including nucleic acid isolation, amplification, and mass spec analysis requires only about 2-3 hours time. Therefore the instant disclosed processes of the invention are faster and less expensive to perform than existing DNA detection systems. In addition, because the instant disclosed processes allow the nucleic acid fragments to be identified and detected at the same time by their specific molecular weights (an unambiguous physical standard), the disclosed processes are also much more accurate and reliable than currently available procedures.

BRIEF DESCRIPTION OF THE FIGURES

[0023] FIG. 1A is a diagram showing a process for performing mass spectrometric analysis on one target detection site (TDS) contained within a target nucleic acid molecule (T), which has been obtained from a biological sample. A specific capture sequence (C) is attached to a solid support (SS) via a spacer (S). The capture sequence is chosen to specifically hybridize with a complementary sequence on the target nucleic acid molecule (T), known as the target capture site (TCS). The spacer (S) facilitates unhindered hybridization. A detector nucleic acid sequence (D), which is complementary to the TDS is then contacted with the TDS. Hybridization between D and the TDS can be detected by mass spectrometry.

[0024] FIG. 1B is a diagram showing a process for performing mass spectrometric analysis on at least one target detection site (here TDS 1 and TDS 2) via direct linkage to a solid support. The target sequence (T) containing the target detection site (TDS 1 and TDS 2) is immobilized to a solid support via the formation of a reversible or irreversible bond formed between an appropriate functionality (L) on the target nucleic acid molecule (T) and an appropriate functionality (L) on the solid support. Detector nucleic acid sequences (here D1 and D2), which are complementary to a target detection site (TDS 1 or TDS 2) are then contacted with the TDS. Hybridization between TDS 1 and D1 and/or TDS 2 and D2 can be detected and distinguished based on molecular weight differences.

[0025] FIG. 1C is a diagram showing a process for detecting a wildtype (D^{wt}) and/or a mutant (D^{mut}) sequence in a target (T) nucleic acid molecule. As in FIG. 1A, a specific capture sequence (C) is attached to a solid support (SS) via a spacer (S). In addition the capture sequence is chosen to specifically interact with a complementary sequence on the target sequence (T), the target capture site (TCS) to be detected through hybridization. However, the target detection site (TDS) includes mutation, X, which changes the molecular weight; mutated target detection sites can be distinguished from wildtype by mass spectrometry. Preferably, the detector nucleic acid molecule (D) is designed so that the mutation is in the middle of the molecule and therefore would not lead to a stable hybrid if the wildtype detector oligonucleotide (D^{wt}) is contacted with the target detector sequence, e.g. as a control. The mutation can also be detected if the mutated detector oligonucleotide (D^{mut}) with the matching base as the mutated position is used for hybridization. If a nucleic acid molecule obtained from a biological sample is heterozygous for the particular sequence (i.e. contain both D^{wt} and D^{mut}) both D^{wt} and D^{mut} will be bound to the appropriate strand and the mass difference allows both D^{wt} and D^{mut} to be detected simultaneously.

[0026] FIG. 2 is a diagram showing a process in which several mutations are simultaneously detected on one target sequence by employing corresponding detector oligonucleotides. The molecular weight differences between the detector oligonucleotides D1, D2 and D3 must be large enough so that simultaneous detection (multiplexing) is possible. This can be achieved either by the sequence itself (composition or length) or by the introduction of mass-modifying functionalities M1-M3 into the detector oligonucleotide.

[0027] FIG. 3 is a diagram showing still another multiplex detection format. In this embodiment, differentiation is accomplished by employing different specific capture sequences which are position-specifically immobilized on a flat surface (e.g., a 'chip array'). If different target sequences T1-Tn are present, their capture sites TCS1-TCSn will interact with complementary immobilized capture sequences C1-Cn. Detection is achieved by employing appropriately mass differentiated detector oligonucleotides D1-Dn, which are mass differentiated either by their sequences or by mass modifying functionalities M1-Mn.

[0028] FIG. 4 is a diagram showing a format wherein a predesigned target capture site (TCS) is incorporated into the target sequence using PCR amplification. Only one strand is captured, the other is removed (e.g., based on the interaction between biotin and streptavidin coated magnetic beads). If the biotin is attached to primer 1 the other strand can be appropriately marked by a TCS. Detection is as described above through the interaction of a specific detector oligonucleotide D with the corresponding target detection site TDS via mass spectrometry.

[0029] FIG. 5 is a diagram showing how amplification (here ligation chain reaction (LCR)) products can be prepared and detected by mass spectrometry. Mass differentiation can be achieved by the mass modifying functionalities (M1 and M2) attached to primers (P1 and P4 respectively). Detection by mass spectrometry can be accomplished directly (i.e.) without employing immobilization and target capturing sites (TCS). Multiple LCR reaction can be performed in parallel by providing an ordered array of capturing sequences (C). This format allows separation of the ligation products and spot by spot identification via mass spectrometry or multiplexing if mass differentiation is sufficient.

[0030] FIG. 6A is a diagram showing mass spectrometric analysis of a nucleic acid molecule, which has been amplified by a transcription amplification procedure. An RNA sequence is captured via its TCS sequence, so that wildtype and mutated target detection sites can be detected as above by employing appropriate detector oligonucleotides (D).

[0031] FIG. 6B is a diagram showing multiplexing to detect two different (mutated) sites on the same RNA in a simultaneous fashion using mass-modified detector oligonucleotides M1-D1 and M2-D2.

[0032] FIG. 6C is a diagram of a different multiplexing procedure for detection of specific mutations by employing mass modified dideoxynucleoside or 3'-deoxynucleoside triphosphates and an RNA dependent DNA polymerase. Alternatively, DNA dependent RNA polymerase and ribonucleotide triphosphates can be employed. This format allows for simultaneous detection of all four base possibilities at the site of a mutation (X).

[0033] FIG. 7A is a diagram showing a process for performing mass spectrometric analysis on one target detec-

tion site (TDS) contained within a target nucleic acid molecule (T), which has been obtained from a biological sample. A specific capture sequence (C) is attached to a solid support (SS) via a spacer (S). The capture sequence is chosen to specifically hybridize with a complementary sequence on T known as the target capture site (TCS). A nucleic acid molecule that is complementary to a portion of the TDS is hybridized to the TDS 5' of the site of a mutation (X) within the TDS. The addition of a complete set of dideoxynucleosides or 3'-deoxynucleoside triphosphates (e.g. pppAdd, pppTdd, pppCdd and pppGdd) and a DNA dependent DNA polymerase allows for the addition only of the one dideoxynucleoside or 3'-deoxynucleoside triphosphate that is complementary to X.

[0034] FIG. 7B is a diagram showing a process for performing mass spectrometric analysis to determine the presence of a mutation at a potential mutation site (M) within a nucleic acid molecule. This format allows for simultaneous analysis of both alleles (A) and (B) of a double stranded target nucleic acid molecule, so that a diagnosis of homozygous normal, homozygous mutant or heterozygous can be provided. Allele A and B are each hybridized with complementary oligonucleotides ((C) and (D) respectively), that hybridize to A and B within a region that includes M. Each heteroduplex is then contacted with a single strand specific endonuclease, so that a mismatch at M, indicating the presence of a mutation, results in the cleavage of (C) and/or (D), which can then be detected by mass spectrometry.

[0035] FIG. 8 is a diagram showing how both strands of a target DNA can be prepared for detection using transcription vectors having two different promoters at opposite locations (e.g. the SP 6 and T7 promoter). This format is particularly useful for detecting heterozygous target detection sites (TDS). Employing the SP 6 or the T7 RNA polymerase both strands could be transcribed separately or simultaneously. Both RNAs can be specifically captured and simultaneously detected using appropriately mass-differentiated detector oligonucleotides. This can be accomplished either directly in solution or by parallel processing of many target sequences on an ordered array of specifically immobilized capturing sequences.

[0036] FIG. 9 is a diagram showing how RNA prepared as described in FIGS. 6, 7 and 8 can be specifically digested using one or more ribonucleases and the fragments captured on a solid support carrying the corresponding complementary sequences. Hybridization events and the actual molecular weights of the captured target sequences provide information on whether and where mutations in the gene are present. The array can be analyzed spot by spot using mass spectrometry. DNA can be similarly digested using a cocktail of nucleases including restriction endonucleases. Mutations can be detected by different molecular weights of specific, individual fragments compared to the molecular weights of the wildtype fragments.

[0037] FIG. 10A shows a spectra resulting from the experiment described in the following Example 1. **FIG. 10A-1** shows the absorbance or the 26-mer before hybridization. **FIG. 10A-2** shows the filtrate of the centrifugation after hybridization. **FIG. 10A-3** shows the results after the first wash with 50 mM ammonium citrate. **FIG. 10A-4** shows the results after the second wash with 50 mM ammonium citrate.

[0038] FIG. 10B shows a spectra resulting from the experiment described in the following Example 1 after three washing/centrifugation steps.

[0039] FIG. 10C shows a spectra resulting from the experiment described in the following Example 1 showing the successful desorption of the hybridized 26-mer off of beads.

[0040] FIG. 11 shows a spectra resulting from the experiment described in the following Example 1 showing the successful desorption of the hybridized 40-mer. The efficiency of detection suggests that fragments much longer 40-mers can also be desorbed.

[0041] FIGS. 12A-12C show spectra resulting from the experiment described in the following Example 2 showing the successful desorption and differentiation of an 18-mer and 19-mer by electrospray mass spectrometry: the mixture (12A), peaks resulting from the 18-mer, emphasized (12B) and peaks resulting from the 19-mer, emphasized (12C).

[0042] FIG. 13 is a graphic representation of the process for detecting the Cystic Fibrosis mutation $\Delta F508$ as described in Example 3; N indicates normal and M indicates the mutation detection primer or extended primer.

[0043] FIG. 14 is a mass spectrum of the DNA extension product of a $\Delta F508$ homozygous normal.

[0044] FIG. 15 is a mass spectrum of the DNA extension product of a $\Delta F508$ heterozygous mutant.

[0045] FIG. 16 is a mass spectrum of the DNA extension product of a $\Delta F508$ homozygous normal.

[0046] FIG. 17 is a mass spectrum of the DNA extension product of a $\Delta F508$ homozygous mutant.

[0047] FIG. 18 is mass spectrum of the DNA extension product of a $\Delta F508$ heterozygous mutant.

[0048] FIG. 19 is a graphic representation of various processes for performing apolipoprotein E genotyping.

[0049] FIGS. 20A-20B show the nucleic acid sequence of normal apolipoprotein E (encoded by the E3 allele) and other isotypes encoded by the E2 and E4 alleles.

[0050] FIG. 21A shows the composite restriction pattern for various genotypes of apolipoprotein E.

[0051] FIG. 21B shows the restriction pattern obtained in a 3.5% Metaphor Agarose Gel for various genotypes of apolipoprotein E.

[0052] FIG. 21 C shows the restriction pattern obtained in a 12% polyacrylamide gel for various genotypes of apolipoprotein E.

[0053] FIG. 22A is a chart showing the molecular weights of the 91, 83, 72, 48 and 35 base pair fragments obtained by restriction enzyme cleavage of the E2, E3, and E4 alleles of apolipoprotein E.

[0054] FIG. 22B is the mass spectra of the restriction product of a homozygous E4 apolipoprotein E genotype.

[0055] FIG. 23A is the mass spectra of the restriction product of a homozygous E3 apolipoprotein E genotype.

[0056] FIG. 23B is the mass spectra of the restriction product of a E3/E4 apolipoprotein E genotype.

[0057] FIG. 24 is an autoradiograph of a 7.5% polyacrylamide gel in which 10% (5 μ l) of each PCR was loaded. Sample M: pBR322 Alul digested; sample 1: HBV positive in serological analysis; sample 2: also HBV positive; sample 3: without serological analysis but with an increased level of transaminases, indicating liver disease; sample 4: HBV negative; sample 5: HBV positive by serological analysis; sample 6: HBV negative (-) negative control; (+) positive control). Staining was done with ethidium bromide.

[0058] FIG. 25A is a mass spectrum of sample 1, which is HBV positive. The signal at 20754 Da represent HBV related PCR product (67 nucleotides, calculated mass: 20735 Da). The mass signal at 10390 Da represents $[M+2H]^{2+}$ signal (calculated: 10378 Da).

[0059] FIG. 25B is a mass spectrum of sample 3, which is HBV negative corresponding to PCR, serological and dot blot based assays. The PCR product is generated only in trace amounts. Nevertheless, it is unambiguously detected at 20751 Da (calculated: 20735 Da). The mass signal at 10397 Da represents the $[M+2H]^{2+}$ molecule ion (calculated: 10376 Da).

[0060] FIG. 25C is a mass spectrum of sample 4, which is HBV negative, but CMV positive. As expected, no HBV specific signals could be obtained.

[0061] FIG. 26 shows a part of the *E.coli* lacI gene with binding sites of the complementary oligonucleotides used in the ligase chain reaction (LCR). Here the wildtype sequence is displayed. The mutant contains a point mutation at bp 191 which is also the site of ligation (bold). The mutation is a C to T transition (G to A, respectively). This leads to a T-G mismatch with oligo A (and A-C mismatch with oligo B, respectively).

[0062] FIG. 27 is a 7.5% polyacrylamide gel stained with ethidium bromide. M: chain length standard (pUC19 DNA, MspI digested). Lane 1: LCR with wildtype template. Lane 2: LCR with mutant template. Lane 3: (control) LCR without template. The ligation product (50 bp) was only generated in the positive reactive containing wildtype template.

[0063] FIG. 28 is an HPLC chromatogram of two pooled positive LCRs.

[0064] FIG. 29 shows an HPLC chromatogram under the same conditions but using the mutant template. The small signal of the ligation product is due to either template-free ligation of the educts or to a ligation at a (G-T, A-C) mismatch. The 'false positive' signal is significantly lower than the signal of ligation product with wildtype template depicted in FIG. 28. The analysis of ligation educts leads to 'double-peaks' because two of the oligonucleotides are 5'-phosphorylated.

[0065] FIG. 30A shows the complex signal pattern obtained by MALDI-TOF-MS analysis of Pfu DNA-ligase solution. FIG. 30B shows a MALDI-TOF-spectrum of an unpurified LCR. The mass signal 67569 Da probably represents the Pfu DNA ligase.

[0066] FIG. 31A shows a MALDI-TOF spectrum of two pooled positive LCRs. The signal at 7523 Da represents unligated oligo A (calculated: 7521 Da) whereas the signal at 15449 Da represents the ligation product (calculated: 15450 Da). The signal at 3774 Da is the $[M+2H]^{2+}$ signal of oligo A. The signals in the mass range lower than 2000 Da

are due to the matrix ions. The spectrum corresponds to lane 1 in **FIG. 27** and to the chromatogram in **FIG. 28**. **FIG. 31B** shows a spectrum of two pooled negative LCRs (mutant template). The signal at 7517 Da represents oligo A (calculated: 7521 Da).

[0067] **FIG. 32** shows a spectrum obtained from two pooled LCRs in which only salmon sperm DNA was used as a negative control; only oligo A could be detected, as expected.

[0068] **FIG. 33A** shows a spectrum of two pooled positive LCRs. The purification was done with a combination of ultrafiltration and streptavidin DynaBeads as described in the text. The signal at 15448 Da represents the ligation product (calculated: 15450 Da). The signal at 7527 represents oligo A (calculated: 7521 Da). The signals at 3761 Da is the $[M+2H]^{2+}$ signal of oligo A, where as the signal at 5140 Da is the $[M+3H]^{2+}$ signal of the ligation product. **FIG. 33B** shows a spectrum of two pooled negative LCRs (without template). The signal at 7514 Da represents oligo A (calculated: 7521 Da).

[0069] **FIG. 34A** is a schematic representation of the oligo base extension of the mutation detection primer b using ddTTP. **FIG. 34B** is a schematic representation of the oligo base extension of the mutation detection primer b using ddCTP. The theoretical mass calculation is given in parenthesis. The sequence shown is part of the exon 10 of the CFTR gene that bears the most common cystic fibrosis mutation $\Delta F508$ and more rare mutations $\Delta S07$ as well as Ile506Ser.

[0070] **FIG. 35A** is a MALDI-TOF-MS spectra recorded directly from precipitated oligo base extended primers for mutation detection using ddTTP. **FIG. 35B** is a MALDI-TOF-MS spectra recorded directly from precipitated oligo base extended primers for mutation detection using ddCTP. The spectra on the top of each panel (ddTTP or ddCTP, respectively) shows the annealed primer (CF508) without further extension reaction. The template of diagnosis is pointed out below each spectra and the observed/expected molecular mass are written in parenthesis.

[0071] **FIG. 36** shows the portion of the sequence of pRFc1 DNA, which was used as template for PCR amplification of unmodified and 7-deazapurine containing 99-mer and 200-mer nucleic acids as well as the sequences of the 19-mer primers and the two 18-mer reverse primers.

[0072] **FIG. 37** shows the portion of the nucleotide sequence of M13mp18 RFI DNA, which was used for PCR amplification of unmodified and 7-deazapurine containing 103-mer nucleic acids. Also shown are nucleotide sequences of the 17-mer primers used in the PCR.

[0073] **FIG. 38** shows the result of a polyacrylamide gel electrophoresis of PCR products purified and concentrated for MALDI-TOF MS analysis. M: chain length marker, lane 1: 7-deazapurine containing 99-mer PCR product, lane 2: unmodified 99-mer, lane 3: 7-deazapurine containing 103-mer and lane 4: unmodified 103-mer PCR product.

[0074] **FIG. 39:** an autoradiogram of polyacrylamide gel electrophoresis of PCR reactions carried out with 5'- ^{32}P -labeled primers 1 and 4. Lanes 1 and 2: unmodified and 7-deazapurine modified 200-mer (71123 and 39582 counts), lanes 3 and 4: unmodified and 7-deazapurine modified

200-mer (71123 and 39582 counts) and lanes 5 and 6: unmodified and 7-deazapurine modified 99-mer (173216 and 94400 counts).

[0075] **FIG. 40A** shows a MALDI-TOF mass spectrum of the unmodified 103-mer PCR products (sum of twelve single shot spectra). The mean value of the masses calculated for the two single strands (31768 u and 31759 u) is 31763 u. Mass resolution: 18. **FIG. 40B** shows a MALDI-TOF mass spectrum of 7-deazapurine containing 103-mer PCR product (sum of three single shot spectra). The mean value of the masses calculated for the two single strands (31727 u and 31719 u) is 31723 u. Mass resolution: 67.

[0076] **FIGS. 41A1** and **41A2** show a MALDI-TOF mass spectrum of the unmodified 99-mer PCR product (sum of twenty single shot spectra). Values of the masses calculated for the two single strands: 30261 u and 30794 u. **FIGS. 41B1** and **41B2** show a MALDI-TOF mass spectrum of the 7-deazapurine containing 99-mer PCR product (sum of twelve single shot spectra). Values of the masses calculated for the two single strands: 30224 u and 30750 u.

[0077] **FIG. 42A** shows a MALDI-TOF mass spectrum of the unmodified 200-mer PCR product (sum of 30 single shot spectra). The mean value of the masses calculated for the two single strands (61873 u and 61595 u) is 61734 u. Mass resolution: 28. **FIG. 42B** shows a MALDI-TOF mass spectrum of 7-deazapurine containing 200-mer PCR product (sum of 30 single shot spectra). The mean value of the masses calculated for the two single strands (61772 u and 61514 u) is 61643 u. Mass resolution: 39.

[0078] **FIG. 43A** shows a MALDI-TOF mass spectrum of 7-deazapurine containing 100-mer PCR product with ribo-modified primers. The mean value of the masses calculated for the two single strands (30529 u and 31095 u) is 30812 u. **FIG. 43B** shows a MALDI-TOF mass spectrum of the PCR-product after hydrolytic primer-cleavage. The mean value of the masses calculated for the two single strands (25104 u and 25229 u) is 25167 u. The mean value of the cleaved primers (5437 u and 5918 u) is 5677 u.

[0079] **FIGS. 44A-44D** show the MALDI-TOF mass spectrum of the four sequencing ladders obtained from a 39-mer template (SEQ. ID. No. 13), which was immobilized to streptavidin beads via a 3' biotinylation. A 14-mer primer (SEQ. ID. No. 14) was used in the sequencing.

[0080] **FIG. 45** shows a MALDI-TOF mass spectrum of a solid state sequencing of a 78-mer template (SEQ. ID. No. 15), which was immobilized to streptavidin beads via a 3' biotinylation. A 18-mer primer (SEQ. ID. No. 16) and ddGTP were used in the sequencing.

[0081] **FIG. 46** shows a scheme in which duplex DNA probes with single-stranded overhang capture specific DNA templates and also serve as primers for solid state sequencing.

[0082] **FIG. 47A-47D** shows MALDI-TOF mass spectra obtained from a 5' fluorescent labeled 23-mer (SEQ. ID. No. 19) annealed to an 3' biotinylated 18-mer (SEQ. ID. No. 20), leaving a 5-base overhang, which captured a 15-mer template (SEQ. ID. No. 21).

[0083] **FIGS. 48A-48D** show a stacking fluorogram of the same products obtained from the reaction described in **FIG. 35**, but run on a conventional DNA sequencer.

DETAILED DESCRIPTION OF THE INVENTION

[0084] In general, the instant invention provides mass spectrometric processes for detecting a particular nucleic acid sequence in a biological sample. As used herein, the term "biological sample" refers to any material obtained from any living source (e.g. human, animal, plant, bacteria, fungi, protist, virus). For use in the invention, the biological sample should contain a nucleic acid molecule. Examples of appropriate biological samples for use in the instant invention include: solid materials (e.g. tissue, cell pellets, biopsies) and biological fluids (e.g. urine, blood, saliva, amniotic fluid, mouth wash).

[0085] Nucleic acid molecules can be isolated from a particular biological sample using any of a number of procedures, which are well-known in the art, the particular isolation procedure chosen being appropriate for the particular biological sample. For example, freeze-thaw and alkaline lysis procedures can be useful for obtaining nucleic acid molecules from solid materials; heat and alkaline lysis procedures can be useful for obtaining nucleic acid molecules from urine; and proteinase K extraction can be used to obtain nucleic acid from blood (Rolf, A et al. PCR: Clinical Diagnostics and Research, Springer (1994).

[0086] To obtain an appropriate quantity of nucleic acid molecules on which to perform mass spectrometry, amplification may be necessary. Examples of appropriate amplification procedures for use in the invention include: cloning (Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, 1989), polymerase chain reaction (PCR) (C. R. Newton and A. Graham, PCR, BIOS Publishers, 1994); ligase chain reaction (LCR) (Wiedmann, M., et al., *PCR Methods Appl.* Vol. 3, pp. 57-64 (1994); F. Barnay, *Proc. Natl. Acad. Sci USA* 88: 189-93 (1991)); strand displacement amplification (SDA) (G. Terence Walker et al., *Nucleic Acids Res.* 22: 2670-77 (1994)); and variations such as RT-PCR (Higuchi, et al., *Bio/Technology* 11: 1026-1030 (1993)), allele-specific amplification (ASA) and transcription based processes.

[0087] To facilitate mass spectrometric analysis, a nucleic acid molecule containing a nucleic acid sequence to be detected can be immobilized to a solid support. Examples of appropriate solid supports include beads (e.g. silica gel, controlled pore glass, magnetic, Sephadex/Sephacrose, cellulose), flat surfaces or chips (e.g. glass fiber filters, glass surfaces, metal surface (steel, gold, silver, aluminum, copper and silicon), capillaries, plastic (e.g. polyethylene, polypropylene, polyamide, polyvinylidene difluoride membranes or microtiter plates)); or pins or combs made from similar materials comprising beads or flat surfaces or beads placed into pits in flat surfaces such as wafers (e.g. silicon wafers).

[0088] Immobilization can be accomplished, for example, based on hybridization between a capture nucleic acid sequence, which has already been immobilized to the support and a complementary nucleic acid sequence, which is also contained within the nucleic acid molecule containing the nucleic acid sequence to be detected (**FIG. 1A**). So that hybridization between the complementary nucleic acid molecules is not hindered by the support, the capture nucleic acid can include a spacer region of at least about five nucleotides in length between the solid support and the capture nucleic acid sequence. The duplex formed will be

cleaved under the influence of the laser pulse and desorption can be initiated. The solid support-bound base sequence can be presented through natural oligoribo- or oligodeoxyribo-nucleotide as well as analogs (e.g. thio-modified phosphodiester or phosphotriester backbone) or employing oligonucleotide mimetics such as PNA analogs (see e.g. Nielsen et al., *Science*, 254, 1497 (1991)) which render the base sequence less susceptible to enzymatic degradation and hence increases overall stability of the solid support-bound capture base sequence.

[0089] Alternatively, a target detection site can be directly linked to a solid support via a reversible or irreversible bond between an appropriate functionality (L') on the target nucleic acid molecule (T) and an appropriate functionality (L) on the capture molecule (**FIG. 1B**). A reversible linkage can be such that it is cleaved under the conditions of mass spectrometry (i.e., a photocleavable bond such as a charge transfer complex or a labile bond being formed between relatively stable organic radicals). Furthermore, the linkage can be formed with L' being a quaternary ammonium group, in which case, preferably, the surface of the solid support carries negative charges which repel the negatively charged nucleic acid backbone and thus facilitate the desorption required for analysis by a mass spectrometer. Desorption can occur either by the heat created by the laser pulse and/or, depending on L', by specific absorption of laser energy which is in resonance with the L' chromophore.

[0090] By way of example, the L-L' chemistry can be of a type of disulfide bond (chemically cleavable, for example, by mercaptoethanol or dithioerythritol), a biotin/streptavidin system, a heterobifunctional derivative of a trityl ether group (Gildea et al., "A Versatile Acid-Labile Linker for Modification of Synthetic Biomolecules," *Tetrahedron Letters* 31: 7095 (1990)) which can be cleaved under mildly acidic conditions as well as under conditions of mass spectrometry, a levulinyl group cleavable under almost neutral conditions with a hydrazinium/acetate buffer, an arginine-arginine or lysine-lysine bond cleavable by an endopeptidase enzyme like trypsin or a pyrophosphate bond cleavable by a pyrophosphatase, or a ribonucleotide bond in between the oligodeoxynucleotide sequence, which can be cleaved, for example, by a ribonuclease or alkali.

[0091] The functionalities, L and L', can also form a charge transfer complex and thereby form the temporary L-L' linkage. Since in many cases the "charge-transfer band" can be determined by UV/vis spectrometry (see e.g. *Organic Charge Transfer Complex* by R. Foster, Academic Press, 1969), the laser energy can be tuned to the corresponding energy of the charge-transfer wavelength and, thus, a specific desorption off the solid support can be initiated. Those skilled in the art will recognize that several combinations can serve this purpose and that the donor functionality can be either on the solid support or coupled to the nucleic acid molecule to be detected or vice versa.

[0092] In yet another approach, a reversible L-L' linkage can be generated by homolytically forming relatively stable radicals. Under the influence of the laser pulse, desorption (as discussed above) as well as ionization will take place at the radical position. Those skilled in the art will recognize that other organic radicals can be selected and that, in relation to the dissociation energies needed to homolytically cleave the bond between them, a corresponding laser wave-

length can be selected (see e.g., *Reactive Molecules* by C. Wentrup, John Wiley & Sons, 1984).

[0093] An anchoring function L' can also be incorporated into a target capturing sequence (TCS) by using appropriate primers during an amplification procedure, such as PCR (FIG. 4), LCR (FIG. 5) or transcription amplification (FIG. 6A).

[0094] Prior to mass spectrometric analysis, it may be useful to "condition" nucleic acid molecules, for example to decrease the laser energy required for volatilization and/or to minimize fragmentation. Conditioning is preferably performed while a target detection site is immobilized. An example of conditioning is modification of the phosphodiester backbone of the nucleic acid molecule (e.g. cation exchange), which can be useful for eliminating peak broadening due to a heterogeneity in the cations bound per nucleotide unit. Contacting a nucleic acid molecule with an alkylating agent such as alkyl iodide, iodoacetamide, β -iodoethanol, 2,3-epoxy-1-propanol, the monothio phosphodiester bonds of a nucleic acid molecule can be transformed into a phosphotriester bond. Likewise, phosphodiester bonds may be transformed to uncharged derivatives employing trialkylsilyl chlorides. Further conditioning involves incorporating nucleotides which reduce sensitivity for depurination (fragmentation during MS) such as N7- or N9-deazapurine nucleotides, or RNA building blocks or using oligonucleotide triesters or incorporating phosphorothioate functions which are alkylated or employing oligonucleotide mimetics such as PNA.

[0095] For certain applications, it may be useful to simultaneously detect more than one (mutated) loci on a particular captured nucleic acid fragment (on one spot of an array) or it may be useful to perform parallel processing by using oligonucleotide or oligonucleotide mimetic arrays on various solid supports. "Multiplexing" can be achieved by several different methodologies. For example, several mutations can be simultaneously detected on one target sequence by employing corresponding detector (probe) molecules (e.g. oligonucleotides or oligonucleotide mimetics). However, the molecular weight differences between the detector oligonucleotides D1, D2 and D3 must be large enough so that simultaneous detection (multiplexing) is possible. This can be achieved either by the sequence itself (composition or length) or by the introduction of mass-modifying functionalities M1-M3 into the detector oligonucleotide. (FIG. 2)?

[0096] Mass modifying moieties can be attached, for instance, to either the 5'-end of the oligonucleotide (M^1), to the nucleobase (or bases) (M^2 , M^7), to the phosphate backbone (M^3), and to the 2'-position of the nucleoside (nucleosides) (M^4 , M^6) or/and to the terminal 3'-position (M^5). Examples of mass modifying moieties include, for example, a halogen, an azido, or of the type, XR, wherein X is a linking group and R is a mass-modifying functionality. The mass-modifying functionality can thus be used to introduce defined mass increments into the oligonucleotide molecule.

[0097] Here the mass-modifying moiety, M, can be attached either to the nucleobase, M^2 (in case of the c⁷-deazanucleosides also to C-7, M^7), to the triphosphate group at the alpha phosphate, M^3 , or to the 2'-position of the sugar ring of the nucleoside triphosphate, M^4 and M^6 . Furthermore, the mass-modifying functionality can be added so as to affect chain termination, such as by attaching it to the

3'-position of the sugar ring in the nucleoside triphosphate, M^5 . For those skilled in the art, it is clear that many combinations can serve the purpose of the invention equally well. In the same way, those skilled in the art will recognize that chain-elongating nucleoside triphosphates can also be mass-modified in a similar fashion with numerous variations and combinations in functionality and attachment positions.

[0098] Without limiting the scope of the invention, the mass-modification, M, can be introduced for X in XR as well as using oligo-/polyethylene glycol derivatives for R. The mass-modifying increment in this case is 44, i.e. five different mass-modified species can be generated by just changing m from 0 to 4 thus adding mass units of 45 (m=0), 89 (m=1), 133 (m=2), 177 (m=3) and 221 (m=4) to the nucleic acid molecule (e.g. detector oligonucleotide (D) or the nucleoside triphosphates (FIGS. 3 and 6C), respectively). The oligo/polyethylene glycols can also be monoalkylated by a lower alkyl such as methyl, ethyl, propyl, isopropyl, t-butyl and the like. A selection of linking functionalities, X, are also illustrated. Other chemistries can be used in the mass-modified compounds, as for example, those described recently in *Oligonucleotides and Analogues, A Practical Approach* (F. Eckstein, editor, IRL Press, Oxford, 1991).

[0099] In yet another embodiment, various mass-modifying functionalities, R, other than oligo/polyethylene glycols, can be selected and attached via appropriate linking chemistries, X. A simple mass-modification can be achieved by substituting H for halogens like F, Cl, Br and/or I, or pseudohalogens such as SCN, NCS, or by using different alkyl, aryl or aralkyl moieties such as methyl, ethyl, propyl, isopropyl, t-butyl, hexyl, phenyl, substituted phenyl, benzyl, or functional groups such as CH_2F , CHF_2 , CF_3 , $\text{Si}(\text{CH}_3)_3$, $\text{Si}(\text{CH}_3)_2(\text{C}_2\text{H}_5)$, $\text{Si}(\text{CH}_3)(\text{C}_2\text{H}_5)_2$, $\text{Si}(\text{C}_2\text{H}_5)_3$. Yet another mass-modification can be obtained by attaching homo- or heteropeptides through the nucleic acid molecule (e.g. detector (D)) or nucleoside triphosphates. One example useful in generating mass-modified species with a mass increment of 57 is the attachment of oligoglycines, e.g. mass-modifications of 74 (r=1, m=0), 131 (r=1, m=2), 188 (r=1, m=3), 245 (r=1, m=4) are achieved. Simple oligoamides also can be used, e.g., mass-modifications of 74 (r=1, m=0), 88 (r=2, m=0), 102 (r=3, m=0), 116 (r=4, m=0), et obtainable. For those skilled in the art, it will be obvious that there are numerous possibilities in addition to those mentioned above.

[0100] As used herein, the superscript O-i designates i+1 mass differentiated nucleotides, primers or tags. In some instances, the superscript O can designate an unmodified species of a particular reactant, and the superscript i can designate the i-th mass-modified species of that reactant. If, for example, more than one species of nucleic acids are to be concurrently detected, then i+1 different mass-modified detector oligonucleotides (D^0 , D^1 , . . . D^i) can be used to distinguish each species of mass modified detector oligonucleotides (D) from the others by mass spectrometry.

[0101] Different mass-modified detector oligonucleotides can be used to simultaneously detect all possible variants/mutants simultaneously (FIG. 6B). Alternatively, all four base permutations at the site of a mutation can be detected by designing and positioning a detector oligonucleotide, so that it serves as a primer for a DNA/RNA polymerase (FIG. 6C). For example, mass modifications also can be incorporated during the amplification process.

[0102] FIG. 3 shows a different multiplex detection format, in which differentiation is accomplished by employing different specific capture sequences which are position-specifically immobilized on a flat surface (e.g. a 'chip array'). If different target sequences T1-Tn are present, their target capture sites TCS1-TCSn will specifically interact with complementary immobilized capture sequences C1-Cn. Detection is achieved by employing appropriately mass differentiated detector oligonucleotides D1-Dn, which are mass differentiated either by their sequences or by mass modifying functionalities M1-Mn.

[0103] Preferred mass spectrometer formats for use in the invention are matrix assisted laser desorption/ionization (MALDI), electrospray (ES), ion cyclotron resonance (ICR) and Fourier Transform. For ES, the samples, dissolved in water or in a volatile buffer, are injected either continuously or discontinuously into an atmospheric pressure ionization interface (API) and then mass analyzed by a quadrupole. The generation of multiple ion peaks which can be obtained using ES mass spectrometry can increase the accuracy of the mass determination. Even more detailed information on the specific structure can be obtained using an MS/MS quadrupole configuration.

[0104] In MALDI mass spectrometry, various mass analyzers can be used, e.g., magnetic sector/magnetic deflection instruments in single or triple quadrupole mode (MS/MS), Fourier transform and time-of-flight (TOF) configurations as is known in the art of mass spectrometry. For the desorption/ionization process, numerous matrix/laser combinations can be used. Ion-trap and reflectron configurations can also be employed.

[0105] The mass spectrometric processes described above can be used, for example, to diagnose any of the more than 3000 genetic diseases currently known (e.g. hemophilias, thalassemias, Duchenne Muscular Dystrophy (DMD), Huntington's Disease (HD), Alzheimer's Disease and Cystic Fibrosis (CF)) or to be identified.

[0106] The following Example 3 provides a mass spectrometer method for detecting a mutation (Δ F508) of the cystic fibrosis transmembrane conductance regulator gene (CFTR), which differs by only three base pairs (900 daltons) from the wild type of CFTR gene. As described further in Example 3, the detection is based on a single-tube, competitive oligonucleotide single base extension (COSBE) reaction using a pair of primers with the 3'-terminal base complementary to either the normal or mutant allele. Upon hybridization and addition of a polymerase and the nucleoside triphosphate one base downstream, only those primers properly annealed (i.e., no 3'-terminal mismatch) are extended; products are resolved by molecular weight shifts as determined by matrix assisted laser desorption/ionization time-of-flight mass spectrometry. For the cystic fibrosis Δ F508 polymorphism, 28-mer 'normal' (N) and 30-mer 'mutant' (M) primers generate 29- and 31-mers for N and M homozygotes, respectively, and both for heterozygotes. Since primer and product molecular weights are relatively low (<10 kDa) and the mass difference between these are at least that of a single ~300 Da nucleotide unit, low resolution instrumentation is suitable for such measurements.

[0107] In addition to mutated genes, which result in genetic disease, certain birth defects are the result of chromosomal abnormalities such as Trisomy 21 (Down's syn-

drome), Trisomy 13 (Platau Syndrome), Trisomy 18 (Edward's Syndrome), Monosomy X (Turner's Syndrome) and other sex chromosome aneuploidies such as Klinefelter's Syndrome (XXY).

[0108] Further, there is growing evidence that certain DNA sequences may predispose an individual to any of a number of diseases such as diabetes, arteriosclerosis, obesity, various autoimmune diseases and cancer (e.g. colorectal, breast, ovarian, lung); chromosomal abnormality (either prenatally or postnatally); or a predisposition to a disease or condition (e.g. obesity, arteriosclerosis, cancer). Also, the detection of "DNA fingerprints", e.g. polymorphisms, such as "microsatellite sequences", are useful for determining identity or heredity (e.g. paternity or maternity).

[0109] The following Example 4 provides a mass spectrometer method for identifying any of the three different isoforms of human apolipoprotein E, which are coded by the E2, E3 and E4 alleles. Here the molecular weights of DNA fragments obtained after restriction with appropriate restriction endonucleases can be used to detect the presence of a mutation.

[0110] Depending on the biological sample, the diagnosis for a genetic disease, chromosomal aneuploidy or genetic predisposition can be preformed either pre- or post-natally.

[0111] Viruses, bacteria, fungi and other organisms contain distinct nucleic acid sequences, which are different from the sequences contained in the host cell. Detecting or quantitating nucleic acid sequences that are specific to the infectious organism is important for diagnosing or monitoring infection. Examples of disease causing viruses that infect humans and animals and which may be detected by the disclosed processes include: Retroviridae (e.g., human immunodeficiency viruses, such as HIV-1 (also referred to as HTLV-III, LAV or HTLV-III/LAV, see Ratner, L. et al., *Nature* 313: 227-284 (1985); and Wain Hobson, S., et al., *Cell* 40: 9-17 (1985)); HIV-2 (see Guyader et al., *Nature* 328: 662-669 (1987); European Patent Publication No. 0 269 520; Chakraborti et al., *Nature* 328: 543-547 (1987); and European Patent Application No. 0 655 501); and other isolates, such as HIV-LP (International Publication No. WO 94/00562 entitled "A Novel Human Immunodeficiency Virus"; Picornaviridae (e.g., polio viruses, hepatitis A virus, (Gust, I. D., et al., *Intervirology*, Vol. 20, pp. 1-7 (1983); enteroviruses, human coxsackie viruses, rhinoviruses, echoviruses); Calciviridae (e.g., strains that cause gastroenteritis); Togaviridae (e.g., equine encephalitis viruses, rubella viruses); Flaviviridae (e.g., dengue viruses, encephalitis viruses, yellow fever viruses); Coronaviridae (e.g., coronaviruses); Rhabdoviridae (e.g., vesicular stomatitis viruses, rabies viruses); Filoviridae (e.g., ebola viruses); Paramyxoviridae (e.g., parainfluenza viruses, mumps virus, measles virus, respiratory syncytial virus); Orthomyxoviridae (e.g., influenza viruses); Bunyaviridae (e.g., Hantaan viruses, bunya viruses, phleboviruses and Nairo viruses); Arenaviridae (hemorrhagic fever viruses); Reoviridae (e.g., reoviruses, orbiviruses and rotaviruses); Birnaviridae; Hepadnaviridae (Hepatitis B virus); Parvoviridae (parvoviruses); Papovaviridae (papilloma viruses, polyoma viruses); Adenoviridae (most adenoviruses); Herpesviridae (herpes simplex virus (HSV) 1 and 2, varicella zoster virus, cytomegalovirus (CMV), herpes viruses); Poxviridae (variola viruses, vaccinia viruses, pox viruses); and Iridoviridae (e.g., African

swine fever virus); and unclassified viruses (e.g., the etiological agents of Spongiform encephalopathies, the agent of delta hepatitis (thought to be a defective satellite of hepatitis B virus), the agents of non-A, non-B hepatitis (class 1=internally transmitted; class 2=parenterally transmitted (i.e., Hepatitis C); Norwalk and related viruses, and astroviruses).

[0112] Examples of infectious bacteria include: *Helicobacter pylori*, *Borrelia burgdorferi*, *Legionella pneumophila*, *Mycobacteria* sps (e.g. *M. tuberculosis*, *M. avium*, *M. intracellulare*, *M. kansasii*, *M. goodii*), *Staphylococcus aureus*, *Neisseria gonorrhoeae*, *Neisseria meningitidis*, *Listeria monocytogenes*, *Streptococcus pyogenes* (Group A Streptococcus), *Streptococcus agalactiae* (Group B Streptococcus), *Streptococcus (viridans group)*, *Streptococcus faecalis*, *Streptococcus bovis*, *Streptococcus (anaerobic sps.)*, *Streptococcus pneumoniae*, pathogenic *Campylobacter* sp., *Enterococcus* sp., *Haemophilus influenzae*, *Bacillus anthracis*, *Corynebacterium diphtheriae*, *Corynebacterium* sp., *Erysipelothrix rhusiopathiae*, *Clostridium perfringens*, *Clostridium tetani*, *Enterobacter aerogenes*, *Klebsiella pneumoniae*, *Pasteurella multocida*, *Bacteroides* sp., *Fusobacterium nucleatum*, *Streptobacillus moniliformis*, *Treponema pallidum*, *Treponema pertense*, *Leptospira*, and *Actinomyces israelii*.

[0113] Examples of infectious fungi include: *Cryptococcus neoformans*, *Histoplasma capsulatum*, *Coccidioides immitis*, *Blastomyces dermatitidis*, *Chlamydia trachomatis*, *Candida albicans*. Other infectious organisms (i.e., protists) include: *Plasmodium falciparum* and *Toxoplasma gondii*.

[0114] The following Example 5 provides a nested PCR and mass spectrometer based method that was used to detect hepatitis B virus (HBV) DNA in blood samples. Similarly, other blood-borne viruses (e.g., HIV-1, HIV-2, hepatitis C virus (HCV), hepatitis A virus (HAV) and other hepatitis viruses (e.g., non-A-non-B hepatitis, hepatitis G, hepatitis E), cytomegalovirus, and herpes simplex virus (HSV)) can be detected each alone or in combination based on the methods described herein.

[0115] Since the sequence of about 16 nucleotides is specific on statistical grounds (even for a genome as large as the human genome), relatively short nucleic acid sequences can be used to detect normal and defective genes in higher organisms and to detect infectious microorganisms (e.g. bacteria, fungi, protists and yeast) and viruses. DNA sequences can even serve as a fingerprint for detection of different individuals within the same species. (Thompson, J. S. and M. W. Thompson, eds., *Genetics in Medicine*, W. B. Saunders Co., Philadelphia, Pa. (1986).

[0116] One process for detecting a wildtype (D^{wt}) and/or a mutant (D^{mut}) sequence in a target (T) nucleic acid molecule is shown in FIG. 1C. A specific capture sequence (C) is attached to a solid support (ss) via a spacer (S). In addition, the capture sequence is chosen to specifically interact with a complementary sequence on the target sequence (T), the target capture site (TCS) to be detected through hybridization. However, if the target detection site (TDS) includes a mutation, X, which increases or decreases the molecular weight, mutated TDS can be distinguished from wildtype by mass spectrometry. For example, in the case of an adenine base (dA) insertion, the difference in molecular weights between D^{wt} and D^{mut} would be about 314 daltons.

[0117] Preferably, the detector nucleic acid (D) is designed such that the mutation would be in the middle of the molecule and the flanking regions are short enough so that a stable hybrid would not be formed if the wildtype detector oligonucleotide (D^{wt}) is contacted with the mutated target detector sequence as a control. The mutation can also be detected if the mutated detector oligonucleotide (D^{mut}) with the matching base at the mutated position is used for hybridization. If a nucleic acid obtained from a biological sample is heterozygous for the particular sequence (i.e. contain both D^{wt} and D^{mut}), both D^{wt} and D^{mut} will be bound to the appropriate strand and the mass difference allows by D^{wt} and D^{mut} to be detected simultaneously.

[0118] The process of this invention makes use of the known sequence information of the target sequence and known mutation sites, although new mutations can also be detected. For example, as shown in FIG. 8, transcription of a nucleic acid molecule obtained from a biological sample can be specifically digested using one or more nucleases and the fragments captured on a solid support carrying the corresponding complementary nucleic acid sequences. Detection of hybridization and the molecular weights of the captured target sequences provide information on whether and where in a gene a mutation is present. Alternatively, DNA can be cleaved by one or more specific endonucleases to form a mixture of fragments. Comparison of the molecular weights between wildtype and mutant fragment mixtures results in mutation detection.

[0119] The present invention is further illustrated by the following examples which should not be construed as limiting in any way. The contents of all cited references (including literature references, issued patents, published patent applications (including international patent application Publication Number WO 94/16101 and U.S. Pat. No. 5,605,798, entitled DNA Sequencing by Mass Spectrometry by H. Köster; and international patent application Publication Number WO 94/21822 and U.S. Pat. No. 5,622,824, entitled "DNA Sequencing by Mass Spectrometry via Exonuclease Degradation" by H. Köster), and co-pending patent applications, (including U.S. patent application Ser. No. 08/406,199, now U.S. Pat. No. 5,605,798 entitled DNA Diagnostics Based on Mass Spectrometry by H. Köster), as cited throughout this application are hereby expressly incorporated by reference.

EXAMPLE 1

MALDI-TOF Desorption of Oligonucleotides Directly on Solid Supports

[0120] 1 g CPC (Controlled Pore Glass) was functionalized with 3-(triethoxysilyl)-epoxypropan to form OH-groups on the polymer surface. A standard oligonucleotide synthesis with 13 mg of the OH-CPG on a DNA synthesizer (Milligen, Model 7500) employing β -cyanoethyl-phosphoramidites (Sinha et al., *Nucleic Acids Res.* 12: 4539 (1984)) and TAC N-protecting groups (Köster et al., *Tetrahedron* 37: 362 (1981)) was performed to synthesize a 3'-T₅-50-mer oligonucleotide sequence in which 50 nucleotides are complementary to a "hypothetical" 50-mer sequence. T₅ serves as a spacer. Deprotection with saturated ammonia in methanol at room temperature for 2 hours furnished according to the determination of the DMT group CPG which contained about 10 μ mol 55-mer/g CPG. This

55-mer served as a template for hybridizations with a 26-mer (with 5'-DMT group) and a 40-mer (without DMT group). The reaction volume is 100 μ l and contains about 1 nmol CPG bound 55-mer as template, an equimolar amount of oligonucleotide in solution (26-mer or 40-mer) in 20 mM Tris-HCl, pH 7.5, 10 mM MgCl₂ and 25 mM NaCl. The mixture was heated for 10' at 65° C. and cooled at 37° C. during 30' (annealing). The oligonucleotide which has not been hybridized to the polymer-bound template were removed by centrifugation and three subsequent washing/centrifugation steps with 100 μ l each of ice-cold 50 mM ammonium citrate. The beads were air-dried and mixed with matrix solution (3-hydroxypicolinic acid/10 mM ammonium citrate in acetonitrile/water, 1:1), and analyzed by MALDI-TOF mass spectrometry. The results are presented in FIGS. 10A-10C and 11.

EXAMPLE 2

Electrospray (ES) Desorption and Differentiation of an 18-mer and 19-mer

[0121] DNA fragments at a concentration of 50 pmole/ μ l in 2-propanol/10 mM ammonium carbonate (1/9, v/v) were analyzed simultaneously by an electrospray mass spectrometer.

[0122] The successful desorption and differentiation of an 18-mer and 19-mer by electrospray mass spectrometry is shown in FIGS. 12A-12C.

EXAMPLE 3

Detection of the Cystic Fibrosis Mutation, Δ F508, by Single Step Dideoxy Extension and Analysis by MALDI-TOF Mass Spectrometry

Material and Methods

[0123] PCR Amplification and Strand Immobilization. Amplification was carried out with exon 10 specific primers using standard PCR conditions (30 cycles: 1'@95° C., 1'@55° C., 2'@72° C.); the reverse primer was 5' labelled with biotin and column purified (Oligopurification Cartridge, Cruachem). After amplification the PCR products were purified by column separation (Qiagen Quickspin) and immobilized on streptavidin coated magnetic beads (Dyna-Beads, Dynal, Norway) according to their standard protocol; DNA was denatured using 0.1M NaOH and washed with 0.1M NaOH, 1x B+W buffer and TE buffer to remove the non-biotinylated sense strand.

[0124] COSBE Conditions. The beads containing ligated antisense strand were resuspended in 18 μ l of Reaction mix (2 μ l 10xTaq buffer, 1 μ l (1 unit) Taq Polymerase, 2 μ l of 2 mM dGTP, and 13 μ l H₂O) and incubated at 80° C. for 5' before the addition of Reaction mix 2 (100 ng each of COSBE primers). The temperature was reduced to 60° C. and the mixtures incubated for 5' annealing/extension period; the beads were then washed in 25 mM triethylammonium acetate (TEAA) followed by 50 mM ammonium citrate.

[0125] Primer Sequences. All primers were synthesized on a Perspective Biosystems Expedite 8900 DNA Synthesizer using conventional phosphoramidite chemistry (Sinha et al. (1984) *Nucleic Acids Res.* 12: 4539. COSBE primers (both

containing an intentional mismatch one base before the 3'-terminus) were those used in a previous ARMS study (Ferrie et al., (1992) *Am J Hum Genet* 51: 251-262) with the exception that two bases were removed from the 5'-end of the normal:

[0126] Ex 10 PCR (Forward): 5'-BIO-GCA AGT GAA TCC TGA GCG TG-3' (SEQ.ID.No. 1)

[0127] Ex 10 PCR (Reverse): 5'-GTG TGA AGG GTT CAT ATG C-3' (SEQ.ID.No. 2)

[0128] COSBE Δ F508-N 5'-ATC TAT ATT CAT CAT AGG AAA CAC CAC A-3' (28-mer) (SEQ.ID.No. 3)

[0129] COSBE Δ F508-M 5'-GTA TCT ATA TTC ATC ATA GGA AAC ACC ATT-3' (30-mer) (SEQ.ID.No. 4).

[0130] Mass Spectrometry. After washing, beads were resuspended in 1 μ l 18 Mohm/cm H₂O. 300 nL each of matrix (Wu et al. (1993) *Rapid Commun Mass Spectrom* 7: 142-146) solution (0.7 M 3-hydroxypicolinic acid, 0.7 M dibasic ammonium citrate in 1:1 H₂O:CH₃CN) and resuspended beads (Tang et al. (1995) *Rapid Commun Mass Spectrom* 8: 727-730) were mixed on a sample target and allowed to air dry. Up to 20 samples were spotted on a probe target disk for introduction into the source region of an unmodified Thermo Bioanalysis (formerly Finnigan) Visions 2000 MALDI-TOF operated in reflectron mode with 5 and 20 kV on the target and conversion dynode, respectively. Theoretical average molecular weights (M_{calc}) were calculated from atomic compositions. Vendor-provided software was used to determine peak centroids using external calibration; 1.08 Da has been subtracted from these to correct for the charge-carrying proton mass to yield the test Mr(exp) values.

[0131] Scheme. Upon annealing to the bound template, the N and M primers (8508.6 and 9148.0 Da, respectively) are presented with dGTP; only primers with proper Watson-Crick base pairing at the variable (V) position are extended by the polymerase. Thus if V pairs with 3-terminal base of N, N is extended to a 8837.9 Da product (N+1). Likewise, if V is properly matched to the M terminus, M is extended to a 9477.3 Da M+1 product.

Results

[0132] FIGS. 14-18 show the representative mass spectra of COSBE reaction products. Better results were obtained when PCR products were purified before the biotinylated anti-sense strand was bound.

EXAMPLE 4

Differentiation of Human Apolipoprotein E Isoforms by Mass Spectrometry

[0133] Apolipoprotein E (Apo E), a protein component of lipoproteins, plays an essential role in lipid metabolism. For example, it is involved with cholesterol transport, metabolism of lipoprotein particles, immunoregulation and activation of a number of lipolytic enzymes.

[0134] There are common isoforms of human Apo E (coded by E2, E3, and E4 alleles). The most common is the E3 allele. The E2 allele has been shown to decrease the

cholesterol level in plasma and therefore may have a protective effect against the development of atherosclerosis. Finally, the E4 isoform has been correlated with increased levels of cholesterol, conferring predisposition to atherosclerosis. Therefore, the identity of the apo E allele of a particular individual is an important determinant of risk for the development of cardiovascular disease.

[0135] As shown in FIG. 19, a sample of DNA encoding apolipoprotein E can be obtained from a subject, amplified (e.g. via PCR); and the PCR product can be digested using an appropriate enzyme (e.g. CfoI). The restriction digest obtained can then be analyzed by a variety of means. As shown in FIGS. 20A-20B, the three isotypes of apolipoprotein E (E2, E3 and E4 have different nucleic acid sequences and therefore also have distinguishable molecular weight values.

[0136] As shown in FIGS. 21A-21C, different Apolipoprotein E genotypes exhibit different restriction patterns in a 3.5% MetPhor Agarose Gel or 12% polyacrylamide gel. As shown in FIGS. 22A-22B and 23A-23B, the various apolipoprotein E genotypes can also be accurately and rapidly determined by mass spectrometry.

EXAMPLE 5

Detection of Hepatitis B Virus in Serum Samples

Materials and Methods

[0137] Sample Preparation

[0138] Phenol/chloroform extraction of viral DNA and the final ethanol precipitation was done according to standard protocols.

[0139] First PCR:

[0140] Each reaction was performed with 5 μ l of the DNA preparation from serum. 15 pmol of each primer and 2 units Taq DNA polymerase (Perkin Elmer, Weiterstadt, Germany) were used. The final concentration of each dNTP was 200 μ M, and the final volume of the reaction was 50 μ l. 10 \times PCR buffer (Perkin Elmer, Weiterstadt, Germany) contained 100 mM Tris-HCl, pH 8.3, 500 mM KCl, 15 mM MgCl₂, and 0.01% gelatine (w/v).

Primer sequences:

Primer 1:
5'-GCTTTTGGGGCATGGACATTGACCCGTATAA-3' (SEQ ID NO 5)

Primer 2:
5'-CTGACTACTAATTCCTGGATGCTGGGTCT-3' (SEQ ID NO 6)

[0141] Nested PCR:

[0142] Each reaction was performed either with 1 μ l of the first reaction or with a 1:10 dilution of the first PCR as template, respectively. 100 pmol of each primer, 2.5 u Pfu(exo-) DNA polymerase (Stratagene, Heidelberg, Germany), a final concentration of 200 μ M of each dNTPs and 5 μ l 10 \times Pfu buffer (200 mM Tris-HCl, pH 8.75, 100 mM KCl, 100 mM (NH₄)₂SO₄, 20 mM MgSO₄, 1% Triton X-100, and 1 mg/ml BSA, (Stratagene, Heidelberg, Germany) were used in a final volume of 50 μ l. The reactions were performed in a thermocycler (OmniGene, MWG-Biotech, Ebersberg, Germany) using the following program:

92° C. for 1 minute, 60° C. for 1 minute and 72° C. for 1 minute with 20 cycles. The sequences of the HBV oligodeoxynucleotides (purchased HPLC-purified at MWG-Biotech, Ebersberg, Germany) are provided below.

[0143] HBV 13: 5'-TTGCCTGAGTGCAGTATGGT-3' (SEQ.ID.No. 7)

[0144] HBV 15bio: Biotin-5'-AGCTCTATATCGG-GAAGCCCT-3' (SEQ.ID.No. 8)

[0145] Purification of PCR Products:

[0146] For the recording of each spectrum, one PCR, 50 μ l, (performed as described above) was used. Purification was done according to the following procedure: Ultrafiltration was done using Ultrafree-MC filtration units (Millipore, Eschborn, Germany) according to the protocol of the provider with centrifugation at 8000 rpm for 20 minutes. 25 μ l (10 μ g/ μ l) streptavidin Dynabeads (Dyna, Hamburg, Germany) were prepared according to the instructions of the manufacturer and resuspended in 25 μ l of B/W buffer (10 mM Tris-HCl, pH7.5, 1 mM EDTA, 2 M NaCl). This suspension was added to the PCR samples still in the filtration unit and the mixture was incubated with gentle shaking for 15 minutes at ambient temperature. The suspension was transferred in a 1.5 ml Eppendorf tube and the supernatant was removed with the aid of a Magnetic Particle Collector, MPC, (Dyna, Hamburg, Germany). The beads were washed twice with 50 μ l of 0.7 M ammonium citrate solution, pH 8.0 (the supernatant was removed each time using the MPC). Cleavage from the beads can be accomplished by using formamide at 90° C. The supernatant was dried in a speedvac for about an hour and resuspended in 4 μ l of ultrapure water (MilliQ UF plus Millipore, Eschborn, Germany). The preparation was used for MALDI-TOF MS analysis.

[0147] MALDI-TOR MS:

[0148] Half a microliter of the sample was pipetted onto the sample holder, then immediately mixed with 0.5 μ l matrix solution (0.7 M 3-hydroxypicolinic acid, 50% acetonitrile, 70 mM ammonium citrate). This mixture was dried at ambient temperature and introduced into the mass spectrometer. All spectra were taken in positive ion mode using a Finnigan MAT Vision 2000 (Finnigan MAT, Bremen, Germany), equipped with a reflectron 5 keV ion source, 20 keV postacceleration) and a 337 nm nitrogen laser. Calibration was done with a mixture of a 40-mer and a 100-mer. Each sample was measured with different laser energies. In the negative samples, the PCR product was detected neither with less nor with higher laser energies. In the positive samples the PCR product was detected at different places of the sample spot and also with varying laser energies.

Results

[0149] A nested PCR system was used for the detection of HBV DNA in blood samples employing oligonucleotides complementary to the c region of the HBV genome (primer 1: beginning at map position 1763, primer 2 beginning at map position 2032 of the complementary strand) encoding the HBV core antigen (HBV cAg). DNA was isolated from patients serum according to standard protocols. A first PCR was performed with the DNA from these preparations using a first set of primers. If HBV DNA was present in the sample, a DNA fragment of 269 bp was generated.

[0150] In the second reaction, primers which were complementary to a region within the PCR fragment generated in the first PCR were used. If HBV-related PCR products were present in the first PCR, a DNA fragment of 67 bp was generated (see FIG. 25A) in this nested PCR. The usage of a nested PCR system for detection provides a high sensitivity and also serves as a specificity control for the external PCR (Rolfs, A. et al., *PCR: Clinical Diagnostics and Research*, Springer, Heidelberg, 1992). A further advantage is that the amount of fragments generated in the second PCR is high enough to ensure an unproblematic detection although purification losses cannot be avoided.

[0151] The samples were purified using ultrafiltration to remove the primers prior to immobilization on streptavidin DynaBeads. This purification was done because the shorter primer fragments were immobilized in higher yield on the beads due to steric reasons. The immobilization was done directly on the ultrafiltration membrane to avoid substance losses due to unspecific absorption on the membrane. Following immobilization, the beads were washed with ammonium citrate to perform cation exchange (Pieles, U. et al., (1992) *Nucleic Acids Res* 21: 3191-3196). The immobilized DNA was cleaved from the beads using 25% ammonia which allows cleavage of DNA from the beads in a very short time, but does not result in an introduction of sodium cations.

[0152] The nested PCRs and the MALDI TOF analysis were performed without knowing the results of serological analysis. Due to the unknown virus titer, each sample of the first PCR was used undiluted as template and in a 1:10 dilution, respectively.

[0153] Sample 1 was collected from a patient with chronic active HBV infection, was positive in HBs and HBe-antigen tests, but negative in a dot blot analysis. Sample 2 was a serum sample from a patient with an active HBV infection and a massive viremia who was HBV positive in a dot blot analysis. Sample 3 was a denatured serum sample, therefore no serological analysis could be performed but an increased level of transaminases indicating liver disease was detected. In autoradiograph analysis (FIG. 24), the first PCR of this sample was negative. Nevertheless, there was some evidence of HBV infection. This sample is of interest for MALDI-TOF analysis, because it demonstrates that even low-level amounts of PCR products can be detected after the purification procedure. Sample 4 was from a patient who was cured of HBV infection. Samples 5 and 6 were collected from patients with a chronic active HBV infection.

[0154] FIG. 24 shows the results of a PAGE analysis of the nested PCR reaction. A PCR product is clearly revealed in samples 1, 2, 5 and 6. In sample 4 no PCR product was

generated; it is indeed HBV negative, according to the serological analysis. Amplification artifacts are visible in lanes 2, 5 and 6, if non-diluted template was used. These artifacts were not generated if the template was used in a 1:10 dilution. In sample 3, PCR product was only detectable if the template was not diluted. The results of PAGE analysis are in agreement with the data obtained by serological analysis except for sample 3 as discussed above.

[0155] FIG. 25A shows a mass spectrum of a nested PCR product from sample number 1 generated and purified as described above. The signal at 20754 Da represents the single stranded PCR product (calculated: 20735 Da, as the average mass of both strands of the PCR product cleaved from the beads). The mass difference of calculated and obtained mass is 19 Da (0.09%). As shown in FIG. 25A, sample number 1 generated a high amount of PCR product, resulting in an unambiguous detection.

[0156] FIG. 25B shows a spectrum obtained from sample number 3. As depicted in FIG. 24, the amount of PCR product generated in this section is significantly lower than that from sample number 1. Nevertheless, the PCR product is clearly revealed with a mass of 20751 Da (calculated 20735). The mass difference is 16 Da (0.08%). The spectrum depicted in FIG. 25C was obtained from sample number 4 which is HBV negative (as is also shown in FIG. 24). As expected no signals corresponding to the PCR product could be detected. All samples shown in FIGS. 25A-25C were analyzed with MALDI-TOF MS, whereby PCR product was detected in all HBV positive samples, but not in the HBV negative samples. These results were reproduced in several independent experiments.

EXAMPLE 6

Analysis of Ligase Chain Reaction Products Via MALDI-TOF Mass Spectrometry

Materials and Methods

[0157] Oligodeoxynucleotides

[0158] Except for the biotinylated oligonucleotide, all other oligonucleotides were synthesized in a 0.2 μ mol scale on a MilliGen 7500 DNA Synthesizer (Millipore, Bedford, Mass., USA) using the β -cyanoethylphosphoramidite method (Sinha, N. D. et al., (1984) *Nucleic Acids Res.* 12: 4539-4577). The oligodeoxynucleotides were RP-HPLC-purified and deprotected according to standard protocols. The biotinylated oligodeoxynucleotide was purchased (HPLC-purified) from Biometra, Göttingen, Germany).

[0159] Sequences and calculated masses of the oligonucleotides used:

Oligodeoxynucleotide A:	5'-p-TTGTGCCACGCGGTGGGAATGTA-3'	(SEQ.ID.No.9)
(7521 Da)		
Oligodeoxynucleotide B:	5'-p-AGCAACGACTGTTGCCCGCCAGTTG-3'	(SEQ.ID.No.10)
(7948 Da)		
Oligodeoxynucleotide C:	5'-bio-TACATTCCCAACCGGTGGCACAAC-3'	(SEQ.ID.No.11)
(7960 Da)		

-continued

Oligodeoxynucleotide D: 5'-p-AACTGGCGGGCAAACAGTCGTTGCT-3' (SEQ.ID.No.12)
(7708 Da)

[0160] 5'-Phosphorylation of Oligonucleotides A and D

[0161] This was performed with polynucleotide Kinase (Boehringer, Mannheim, Germany) according to published procedures; the 5'-phosphorylated oligonucleotides were used unpurified for LCR.

[0162] Ligase Chain Reaction

[0163] The LCR was performed with Pfu DNA ligase and a ligase chain reaction kit (Stratagene, Heidelberg, Germany) containing two different pBluescript KII phagemids, one carrying the wildtype form of the *E. coli* lacI gene and the other one a mutant of the gene with a single point mutation of bp 191 of the lacI gene.

[0164] The following LCR conditions were used for each reaction: 100 pg template DNA (0.74 fmol) with 500 pg sonified salmon sperm DNA as carrier, 25 ng (3.3 pmol) of each 5'-phosphorylated oligonucleotide, 20 ng (2.5 pmol) of each non-phosphorylated oligonucleotide, and 4 u Pfu DNA ligase in a final volume of 20 μ l buffered by Pfu DNA ligase reaction buffer (Stratagene, Heidelberg, Germany). In a model experiment, a chemically synthesized ss 50-mer was used (1 fmol) as template, and in this case oligo C was also biotinylated. All reactions were performed in a thermocycler (OmniGene, MWG-Biotech, Ebersberg, Germany) with the following program: 4 minutes 92° C., 2 minutes 60° C. and 25 cycles of 20 seconds 92° C., 40 seconds 60° C. Except for HPLC analysis, the biotinylated ligation educt C was used. In a control experiment the biotinylated and non-biotinylated oligonucleotides revealed the same gel electrophoretic results. The reactions were analyzed on 7.5% polyacrylamide gels. Ligation product 1 (oligo A and B) calculated mass: 15450 Da; ligation product 2 (oligo C and D) calculated mass: 15387 Da.

[0165] SMART-HPLC

[0166] Ion exchange HPLC (IE HPLC) was performed on the SMART-system (Pharmacia, Freiburg, Germany) using a Pharmacia Mono Q, PC 1.6/5 column. Eluents were buffer A (25 mM Tris-HCl, 1 mM EDTA and 0.3 M NaCl at pH 8.0) and buffer B (same as A, but 1 M NaCl). Starting with 100% A for 5 minutes at a flow rate of 50 μ l/min a gradient was applied from 0 to 70% B in 30 minutes, then increased to 100% B in 2 minutes and held at 100% B for 5 minutes. Two pooled LCR volumes (40 μ l) performed with either wildtype or mutant template were injected.

[0167] Sample Preparation for MALDI-TOF-MS

[0168] Preparation of immobilized DNA: For the recording of each spectrum two LCRs (performed as described above) were pooled and diluted 1:1 with 2x B/W buffer (10 mM Tris-HCl, pH 7.5, 1 mM EDTA, 2 M NaCl). To the samples 5 μ l streptavidin DynaBeads (Dyna, Hamburg, Germany) were added and the mixture was allowed to bind with gentle shaking for 15 minutes at ambient temperature. The supernatant was removed using a Magnetic Particle Collector, MPC (Dyna, Hamburg, Germany) and the beads were washed twice with 50 μ l of 0.7 M ammonium citrate solution (pH 8.0) (the supernatant was removed each time

using the MPC). The beads were resuspended in 1 μ l of ultrapure water (MilliQ, Millipore, Bedford, Mass., USA). This suspension was directly used for MALDI-TOF-MS analysis as described below.

[0169] Combination of ultrafiltration and streptavidin DynaBeads: For the recording of spectrum two LCRs (performed as described above) were pooled, diluted 1:1 with 2x B/W buffer and concentrated with a 5000 NMWL Ultrafree-MC filter unit (Millipore, Eschborn, Germany) according to the instructions of the manufacturer. After concentration, the samples were washed with 300 μ l 1x B/W buffer and 5 μ l streptavidin DynaBeads were added. The beads were washed once on the Ultrafree-MC filtration unit with 300 μ l of 1x B/W buffer and processed as described above. The beads were resuspended in 30 to 50 μ l of 1x B/W buffer and transferred in a 1.5 ml Eppendorf tube. The supernatant was removed and the beads were washed twice with 50 μ l of 0.7 M ammonium citrate (pH 8.0). Finally, the beads were washed once with 30 μ l of acetone and resuspended in 1 μ l of ultrapure water. The ligation mixture, after immobilization on the beads, was used for MALDI-TOF-MS analysis as described below.

[0170] MALDI-TOF-MS

[0171] A suspension of streptavidin-coated magnetic beads with the immobilized DNA was pipetted onto the sample holder, then immediately mixed with 0.5 μ l matrix solution (0.7 M 3-hydroxyisobutyric acid in 50% acetonitrile, 70 mM ammonium citrate). This mixture was dried at ambient temperature and introduced into the mass spectrometer. All spectra were taken in positive ion mode using a Finnigan MAT Vision 2000 (Finnigan MAT, Bremen, Germany), equipped with a reflectron (5 ke V ion source, 20 keV postacceleration) and a nitrogen laser (337 nm). For the analysis of Pfu DNA ligase, 0.5 μ l of the solution was mixed on the sample holder with 1 μ l of matrix solution and prepared as described above. For the analysis of unpurified LCRs, 1 μ l of an LCR was mixed with 1 μ l matrix solution.

Results and Discussion

[0172] The *E. coli* lacI gene served as a simple model system to investigate the suitability of MALDI-TOF-MS as a detection method for products generated in ligase chain reactions. This template system contains an *E. coli* lacI wildtype gene in a pBluescript KII phagemid and an *E. coli* lacI gene carrying a single point mutation at bp 191 (C to T transition) in the same phagemid. Four different oligonucleotides were used, which were ligated only if the *E. coli* lacI wildtype gene was present (**FIG. 26**).

[0173] LCR conditions were optimized using Pfu DNA ligase to obtain at least 1 pmol ligation product in each positive reaction. The ligation reactions were analyzed by polyacrylamide gel electrophoresis (PAGE) and HPLC on the SMART system (**FIGS. 27, 28 and 29**). **FIG. 27** shows a PAGE of a positive LCR with wildtype template (lane 1), a negative LCR with mutant template (lane 2) and a negative control which contains enzyme, oligonucleotides and no template. The gel electrophoresis clearly shows that

the ligation product (50 bp) was produced only in the reaction with wildtype template whereas neither the template carrying the point mutation nor the control reaction with salmon sperm DNA generated amplification products. In FIG. 28, HPLC was used to analyze two pooled LCRs with wildtype template performed under the same conditions. The ligation product was clearly revealed. FIG. 29 shows the results of a HPLC in which two pooled negative LCRs with mutant template were analyzed. These chromatograms confirm the data shown in FIG. 27 and the results taken together clearly demonstrate, that the system generates ligation products in a significant amount only if the wildtype template is provided.

[0174] Appropriate control runs were performed to determine retention times of the different compounds involved in the LCR experiments. These include the four oligonucleotides (A, B, C, and D), a synthetic ds 50-mer (with the same sequence as the ligation product), the wildtype template DNA, sonicated salmon sperm DNA and the Pfu DNA ligase in ligation buffer.

[0175] In order to test which purification procedure should be used before a LCR reaction can be analyzed by MALDI-TOF-MS, aliquots of an unpurified LCR (FIG. 30A) and aliquots of the enzyme stock solution (FIG. 30B) were analyzed with MALDI-TOF-MS. It turned out that appropriate sample preparation is absolutely necessary since all signals in the unpurified LCR correspond to signals obtained in the MALDI-TOF-MS analysis of the Pfu DNA ligase. The calculated mass values of oligo A and the ligation product are 7521 Da and 15450 Da, respectively. The data in FIGS. 30A-30B show that the enzyme solution leads to mass signals which do interfere with the expected signals of the ligation educts and products and therefore makes an unambiguous signal assignment impossible. Furthermore, the spectra showed signals that the detergent Tween 20 that is in the enzyme storage buffer influences the crystallization behavior of the analyte/matrix mixture in an unfavorable way.

[0176] In one purification format streptavidin-coated magnetic beads were used. As was shown in a recent paper, the direct desorption of DNA immobilized by Watson-Crick base pairing to a complementary DNA fragment covalently bound to the beads is possible and the non-biotinylated strand will be desorbed exclusively (Tang, K. et al., *Nucleic Acids Res.* 23: 3126-3131 (1995)). This approach in using immobilized ds DNA ensures that only the non-biotinylated strand will be desorbed. If non-immobilized ds DNA is analyzed both strands are desorbed (Tang, K. et al., *Rapid Comm. Mass Spectrom.* 7: 183-186 (1994)) leading to broad signals depending on the mass difference of the two strands. Therefore, employing this system for LCR, only the non-ligated oligonucleotide A, with a calculated mass of 7521 Da, and the ligation product from oligo A and oligo B (calculated mass: 15450 Da) will be desorbed if oligo C is biotinylated at the 5' end and immobilized on streptavidin-coated beads. This results in a simple and unambiguous identification of the LCR educts and products.

[0177] FIG. 31A shows a MALDI-TOF mass spectrum obtained from two pooled LCRs (performed as described above) purified on streptavidin DynaBeads and desorbed directly from the beads, indicating that the purification method used was efficient (compared with FIGS. 30A-30B).

A signal which represents the unligated oligo A and a signal which corresponds to the ligation product could be detected. The agreement between the calculated and the experimentally found mass values is remarkable and allows an unambiguous peak assignment and accurate detection of the ligation product. In contrast, no ligation product but only oligo A could be detected in the spectrum obtained from two pooled LCRs with mutated template (FIG. 31B). The specificity and selectivity of the LCR conditions and the sensitivity of the MALDI-TOF detection is further demonstrated when performing the ligation reaction in the absence of a specific template. FIG. 32 shows a spectrum obtained from two pooled LCRs in which only salmon sperm DNA was used as a negative control; only oligo A could be detected, as expected.

[0178] While the results shown in FIG. 31A can be correlated to lane 1 of the gel in FIG. 27, the spectrum shown in FIG. 31B is equivalent to lane 2 in FIG. 27, and finally also the spectrum in FIG. 32 corresponds to lane 3 in FIG. 27. The results are in congruence with the HPLC analysis presented in FIGS. 28 and 29. While both gel electrophoresis (FIG. 27) and HPLC (FIGS. 28 and 29) reveal either an excess or almost equal amounts of ligation product over ligation educts, the analysis by MALDI-TOF mass spectrometry produces a smaller signal for the ligation product (FIG. 31A).

[0179] The lower intensity of the ligation product signal could be due to different desorption/ionization efficiencies between 24- and a 50-mer. Since the T_m value of a duplex with 50 compared to 24 base pairs is significantly higher, more 24-mer could be desorbed. A reduction in signal intensity can also result from a higher degree of fragmentation in the case of the longer oligonucleotides.

[0180] Regardless of the purification with streptavidin DynaBeads, FIG. 32 reveals traces of Tween 20 in the region around 2000 Da. Substances with a viscous consistency negatively influence the process of crystallization and therefore can be detrimental to mass spectrometer analysis. Tween 20 and also glycerol which are part of enzyme storage buffers therefore should be removed entirely prior to mass spectrometer analysis. For this reason an improved purification procedure which includes an additional ultrafiltration step prior to treatment with DynaBeads was investigated. Indeed, this sample purification resulted in a significant improvement of MALDI-TOF mass spectrometric performance.

[0181] FIGS. 33A-33B show spectra obtained from two pooled positive (33A) and negative (33B) LCRs, respectively. The positive reaction was performed with a chemically synthesized, single strand 50-mer as template with a sequence equivalent to the ligation product of oligo C and D. Oligo C was 5'-biotinylated. Therefore the template was not detected. As expected, only the ligation product of Oligo A and B (calculated mass 15450 Da) could be desorbed from the immobilized and ligated oligo C and D. This newly generated DNA fragment is represented by the mass signal of 15448 Da in FIG. 33A. Compared to FIG. 31A, this spectrum clearly shows that this method of sample preparation produces signals with improved resolution and intensity.

EXAMPLE 7

Mutation Detection by Solid Phase Oligo Base
Extension of a Primer and Analysis by
MALDI-TOF Mass Spectrometry

Summary

[0182] The solid-phase oligo base extension method detects point mutations and small deletions as well as small insertions in amplified DNA. The method is based on the extension of a detection primer that anneals adjacent to a variable nucleotide position on an affinity-captured amplified template, using a DNA polymerase, a mixture of three dNTPs, and the missing one dideoxy nucleotide. The resulting products are evaluated and resolved by MALDI-TOF mass spectrometry without further labeling procedures. The aim of the following experiment was to determine mutant and wildtype alleles in a fast and reliable manner.

Description of the Experiment

[0183] The method used a single detection primer followed by an oligo-nucleotide extension step to give products, differing in length by some bases specific for mutant or wildtype alleles which can be easily resolved by MALDI-TOF mass spectrometry. The method is described by using as example the exon 10 of the CFTR-gene. Exon 10 of this gene leads, in the homozygous state, to the clinical phenotype of cystic fibrosis.

Materials and Methods

[0184] Genomic DNA

[0185] Genomic DNA were obtained from healthy individuals, individuals homozygous or heterozygous for the $\Delta F508$ mutation, and one individual heterozygous for the 1506S mutation. The wildtype and mutant alleles were confirmed by standard Sanger sequencing.

[0186] PCR Amplification of Exon 10 of the CFTR Gene

[0187] The primers for PCR amplification were CFEx10-F (5'-GCAAGTGAATCCTGAGCGTG-3' (SEQ.ID.No. 13) located in intron 9 and biotinylated) and CFEx10-R (5'-GTGTGAAGGGCGTG-3' (SEQ.ID.No. 14) located in intron 10). Primers were used at a concentration of 8 pmol. Taq-polymerase including 10xbuffer was purchased from Boehringer-Mannheim and dNTPs were obtained from Pharmacia. The total reaction volume was 50 μ l. Cycling conditions for PCR were initially 5 min. at 95° C., followed by 1 min. at 94° C., 45 sec at 53° C., and 30 sec at 72° C. for 40 cycles with a final extension time of 5 min at 72° C.

[0188] Purification of the PCR Products

[0189] Amplification products were purified by using Qiagen's PCR purification kit (No. 28106) according to manufacturer's instructions. The elution of the purified products from the column was done in 50 μ l TE-buffer (10 mM Tris, 1 mM EDTA, pH 7.5).

[0190] Affinity-Capture and Denaturation of the Double Stranded DNA

[0191] 10 μ l aliquots of the purified PCR product were transferred to one well of a streptavidin-coated microtiter plate (No. 1645684 Boehringer-Mannheim or No. 95029262 Labystems). Subsequently, 10 μ l incubation buffer (80 mM sodium phosphate, 400 mM NaCl, 0.4% Tween 20, pH 7.5) and 30 μ l water were added. After incubation for 1 hour at

room temperature the wells were washed three times with 200 μ l washing buffer (40 mM Tris, 1 mM EDTA, 50 mM NaCl, 0.1% Tween 20, pH 8.8). To denature the double stranded DNA, the wells were treated with 100 μ l of a 50 mM NaOH solution for 3 min. Hence, the wells were washed three times with 200 μ l washing buffer.

[0192] Oligo Base Extension Reaction

[0193] The annealing of 25 pmol detection primer (CF508: 5'-CTATATTTCATCATAGGAAACACCA-3' (SEQ ID No. 15)) was performed in 50 μ l annealing buffer (20 mM Tris, 10 mM KCl, 10 mM (NH₄)₂SO₄, 2 mM MgSO₄, 1% Triton X-100, pH 8.75) at 50° C. for 10 min. The wells were washed three times with 200 μ l washing buffer and once in 200 μ l TE buffer. The extension reaction was performed by using some components of the DNA sequencing kit from USB (No. 70770) and dNTPs or ddNTPs from Pharmacia. The total reaction volume was 45 μ l, consisting of 21 μ l sample, 6 μ l Sequenase-buffer, 3 μ l 10 mM DTT solution, 4.5 μ l of 0.5 mM of three dNTPs, 4.5 μ l of 2 mM of the missing one ddNTP, 5.5 μ l glycerol enzyme dilution buffer, 0.25 μ l Sequenase 2.0, and 0.25 μ l pyrophosphatase. The reaction was pipetted on ice and then incubated for 15 min. at room temperature and for 5 min. at 37° C. Hence, the wells were washed three times with 200 μ l washing buffer and once with 60 μ l of a 70 mM NH₄-citrate solution.

[0194] Denaturation and Precipitation of the Extended Primer

[0195] The extended primer was denatured in 50 μ l 10% DMSO (dimethylsulfoxide) in water at 80° C. for 10 min. For precipitation, 10 μ l NH₄-acetate (pH 6.5), 0.5 μ l glycogen (10 mg/ml water, Sigma No. G1765), and 100 μ l absolute ethanol were added to the supernatant and incubated for 1 hour at room temperature. After centrifugation at 13,000 g for 10 min., the pellet was washed in 70% ethanol and resuspended in 1 μ l 18 Mohn/cm H₂O water.

[0196] Sample Preparation and Analysis on MALDI-TOF Mass Spectrometry

[0197] Sample preparation was performed by mixing 0.3 μ l each of matrix solution (0.7 M 3-hydroxypicolinic acid, 0.07 M dibasic ammonium citrate in 1:1 H₂O:CH₃CN) and of resuspended DNA/glycogen pellet on a sample target and allowed to air dry. Up to 20 samples were spotted on a probe target disk for introduction into the source region of an unmodified Thermo Bioanalysis (formerly Finnigan) Visions 2000 MALDI-TOF operated in reflectron mode with 5 and 20 kV on the target and conversion dynode, respectively. Theoretical average molecular mass (M_r (calc)) were calculated from atomic compositions; reported experimental M_r (M_r (exp)) values are those of the singly-protonated form, determined using external calibration.

Results

[0198] The aim of the experiment was to develop a fast and reliable method independent of exact stringencies for mutation detection that leads to high quality and high throughput in the diagnosis of genetic diseases. Therefore a special kind of DNA sequencing (oligo base extension of one mutation detection primer) was combined with the evaluation of the resulting mini-sequencing products by matrix-assisted laser desorption ionization (MALDI) mass spectrometry (MS). The time-of-flight (TOF) reflectron arrangement was chosen as a possible mass measurement system. To prove this hypothesis, the examination was performed with exon 10 of the CFTR-gene, in which some

mutations could lead to the clinical phenotype of cystic fibrosis, the most common monogenetic disease in the Caucasian population.

(Millipore, Bedford, Mass., USA) in 200 nmol scales or purchased from MWG-Biotech (Ebersberg, Germany, primer 3) and Biometra (Goettingen, Germany, primers 6-7).

primer 1: 5'-GTCACCCCTGCAGCTGCAG-3'; (SEQ.ID.No.16)
 primer 2: 5'-TTCTAAAACGACGGCCACT-3'; (SEQ.ID.No.17)
 primer 3: 5'-CTTCACCCGCGATGTTGA-3'; (SEQ.ID.No.18)
 primer 4: 5'-CAGGAAACAGCTATGAC-3'; (SEQ.ID.No.19)
 primer 5: 5'-GTAACACGACGGCCAGT-3'; (SEQ.ID.No.20)
 primer 6: 5'-GTCACCCCTGCAGCTGCAGc-3' (g: RiboG); (SEQ.ID.No.21)
 primer 7: 5'-GTTGTAAAACGAGGGCCAgT-3' (g: RiboG). (SEQ.ID.No.22)

[0199] The schematic presentations given in FIGS. 34A-34B show the expected short sequencing products with the theoretically calculated molecular mass of the wildtype and various mutations of exon 10 of the CFTR-gene. The short sequencing products were produced using either ddTTP (FIG. 34A) or ddCTP (FIG. 34B) to introduce a definitive sequence related stop in the nascent DNA strand. The MALDI-TOF-MS spectra of healthy, mutation heterozygous, and mutation homozygous individuals are presented in FIGS. 35A-35B. All samples were confirmed by standard Sanger sequencing which showed no discrepancy in comparison to the mass spec analysis. The accuracy of the experimental measurements of the various molecular masses was within a range of minus 21.8 and plus 87.1 dalton (Da) to the range expected. This is a definitive interpretation of the results allowed in each case. A further advantage of this procedure is the unambiguous detection of the Δ1507 mutation. In the ddTTP reaction, the wildtype allele would be detected, whereas in the ddCTP reaction the three base pair deletion would be disclosed.

[0200] The method described is highly suitable for the detection of single point mutations or microlesions of DNA. Careful choice of the mutation detection primers will open the window of multiplexing and lead to a high throughput including high quality in genetic diagnosis without any need for exact stringencies necessary in comparable allele-specific procedures. Because of the uniqueness of the genetic information, the oligo base extension of mutation detection primer is applicable in each disease gene or polymorphic region in the genome like variable number of tandem repeats (VNTR) or other single nucleotide polymorphisms (e.g., apolipoprotein E gene).

EXAMPLE 8

Detection of Polymerase Chain Reaction Products
 Containing 7-Deazapurine Moieties With
 Matrix-Assisted Laser Desorption/Ionization
 Time-of-Flight (MALDI-TOF) Mass Spectrometry

Materials and Methods

[0201] PCR Amplifications

[0202] The following oligodeoxynucleotide primers were either synthesized according to standard phosphoramidite chemistry (Sinha, N. D. et al., *Tetrahedron Let.* 24: 5843-5846 (1983); Sinha, N. D. et al., *Nucleic Acids Res.* 12: 4539-4557 (1984)) on a Milligen 7500 DNA synthesizer

[0203] The 99-mer and 200-mer DNA strands (modified and unmodified) as well as the ribo- and 7-deaza-modified 100-mer were amplified from pRFc1 DNA (10 ng, generously supplied S. Feyerabend, University of Hamburg) in 100 μl reaction volume containing 10 mmol/L KCl, 10 mmol/L (NH₄)₂SO₄, 20 mmol/L Tris HCl (pH=8.8), 2 mmol/L MgSO₄, (exo(-))Pseudococcus furiosus (Pfu)-Buffer, Pharmacia, Freiburg, Germany), 0.2 mmol/L each dNTP (Pharmacia, Freiburg, Germany), 1 μl of each primer and 1 unit of exo(-)Pfu DNA polymerase (Stratagene, Heidelberg, Germany).

[0204] For the 99-mer primers 1 and 2, for the 200-mer primers 1 and 3 and for the 100-mer primers 6 and 7 were used. To obtain 7-deazapurine modified nucleic acids, during PCR-amplification dATP and dGTP were replaced with 7-deaza-dATP and 7-deaza-dGTP. The reaction was performed in a thermal cycler (OmniGene, MWG-Biotech, Ebersberg, Germany) using the cycle: denaturation at 95° C. for 1 min., annealing at 51° C. for 1 min. and extension at 72° C. for 1 min. For all PCRs the number of reaction cycles was 30. The reaction was allowed to extend for additional 10 min. at 72° C. after the last cycle.

[0205] The 103-mer DNA strands (modified and unmodified) were amplified from M13mp18 RFI DNA (100 ng, Pharmacia, Freiburg, Germany) in 100 μl reaction volume using primers 4 and 5; all other concentrations were unchanged. The reaction was performed using the cycle: denaturation at 95° C. for 1 min., annealing at 40° C. for 1 min., and extension at 72° C. for 1 min. After 30 cycles for the unmodified and 40 cycles for the modified 103-mer respectively, the samples were incubated for an additional 10 min. at 72° C.

[0206] Synthesis of 5'-[³²P]-Labeled PCR-Primers

[0207] Primers 1 and 4 were 5'-[³²P]-labeled employing T4-polynucleotide kinase (Epicentre Technologies) and (γ-³²P)-ATP (BLU/NGG/502A, Dupont, Germany) according to the protocols of the manufacturer. The reactions were performed substituting 10% of primer 1 and 4 in PCR with the labeled primers under otherwise unchanged reaction conditions. The amplified DNAs were separated by gel electrophoresis on a 10% polyacrylamide gel. The appropriate bands were excised and counted on a Packard TRI-CARB 460C liquid scintillation system (Packard, Conn., USA).

[0208] Primer-Cleavage From Ribo-Modified PCR-Product

[0209] The amplified DNA was purified using Ultrafree-MC filter units (30,000 NMWL), it was then redissolved in 100 μ L of 0.2 mol/L NaOH and heated at 95° C. for 25 minutes. The solution was then acidified with HCl (1 mol/L) and further purified for MALDI-TOF analysis employing Ultrafree-MC filter units (30,000 NMWL) as described below.

[0210] Purification of PCR Products

[0211] All samples were purified and concentrated using Ultrafree-MC units 30,000 NMWL (Millipore, Eschborn, Germany) according to the manufacturer's description. After lyophilization, PCR products were redissolved in 5 μ L (3 μ L for the 200-mer) of ultrapure water. This analyte solution was directly used for MALDI-TOF measurements.

[0212] MALDI-TOF MS

[0213] Aliquots of 0.5 μ L of analyte solution and 0.5 μ L of matrix solution (0.7 mol/L 3-HPA and 0.07 mol/L ammonium citrate in acetonitrile/water (1:1, v/v) were mixed on a flat metallic sample support. After drying at ambient temperature the sample was introduced into the mass spectrometer for analysis. The MALDI-TOF mass spectrometer used was a Finnigan MAT Vision 2000 (Finnigan MAT, Bremen, Germany). Spectra were recorded in the positive ion reflection mode with a 5 keV ion source and 20 keV postacceleration. The instrument was equipped with a nitrogen laser (337 nm wavelength). The vacuum of the system was $3.4 \cdot 10^{-8}$ hPa in the analyzer region and $1.4 \cdot 10^{-7}$ hPa in the source region. Spectra of modified and unmodified DNA samples were obtained with the same relative laser power; external calibration was performed with a mixture of synthetic oligodeoxynucleotides (7- to 50-mer).

Results and Discussion

[0214] Enzymatic Synthesis of 7-Deazapurine-Nucleotide-Containing Nucleic Acids by PCR

[0215] In order to demonstrate the feasibility of MALDI-TOF MS for the rapid, gel-free analysis of short PCR products and to investigate the effect of 7-deazapurine modification of nucleic acids under MALDI-TOF conditions, two different primer-template systems were used to synthesize DNA fragments. Sequences are displayed in FIGS. 36 and 37. While the two single strands of the 103-mer PCR product had nearly equal masses ($\Delta M=8$ u), the two single strands of the 99-mer differed by 526 u.

[0216] Considering that 7-deaza purine nucleotide building blocks for chemical DNA synthesis are approximately 160 times more expensive than regular ones (Product Information, Glen Research Corporation, Sterling, Va.) and their application in standard β -cyano-phosphoramidite chemistry is not trivial (Product Information, Glen Research Corporation, Sterling, Va.; Schneider, K. and B. T. Chait, *Nucleic Acids Res.* 23: 1570 (1995)) the cost of 7-deaza-purine-modified primers would be very high. Therefore, to increase the applicability and scope of the method, all PCRs were performed using unmodified oligonucleotide primers which are routinely available. Substituting dATP and dGTP by c^7 -dATP and c^7 -dGTP in polymerase chain reaction led to products containing approximately 80% 7-deaza-purine-

modified nucleosides for the 99-mer and 103-mer; and about 90% for the 200-mer, respectively. Table I shows the base composition of all PCR products.

TABLE I

Base composition of the 99-mer, 103-mer and 200-mer PCR amplification products (unmodified and 7-deaza purine modified)

DNA-fragments ¹	C	T	A	G	c^7 -deaza-A	c^7 -deaza-G	rel. mod. ²
200-mer s	54	34	56	56	—	—	—
modified 200-mer s	54	34	6	5	50	51	90%
200-mer a	56	56	34	54	—	—	—
modified 200-mer a	56	56	3	4	31	50	92%
103-mer s	28	23	24	28	—	—	—
modified 103-mer s	28	23	6	5	18	23	79%
103-mer a	28	24	23	28	—	—	—
modified 103-mer a	28	24	7	4	16	24	78%
99-mer s	34	21	24	20	—	—	—
modified 99-mer s	34	21	6	5	18	15	75%
99-mer a	20	24	21	34	—	—	—
modified 99-mer a	20	24	3	4	18	30	87%

¹"s" and "a" describe "sense" and "antisense" strands of the double-stranded amplified product.

²indicates relative modification as percentage of 7-deaza-purine-modified nucleotides of total amount of Purine nucleotides.

[0217] However, it remained to be determined whether 80-90% 7-deazapurine modification is sufficient for accurate mass spectrometer detection. It was therefore important to determine whether all purine nucleotides could be substituted during the enzymatic amplification step. This was not trivial since it had been shown that c^7 -dATP cannot fully replace dATP in PCR if Taq DNA polymerase is employed (Seela, F. and A. Roelling, *Nucleic Acids Res.* 20: 55-61 (1992)). Fortunately we found that exo(-)Pfu DNA polymerase indeed could accept c^7 -dATP and c^7 -dGTP in the absence of unmodified purine triphosphates. However, the incorporation was less efficient leading to a lower yield of PCR product (FIG. 38). Ethidium-bromide stains by intercalation with the stacked bases of the DNA-doublestrand. Therefore lower band intensities in the ethidium-bromide stained gel might be artifacts since the modified DNA-strands do not necessarily need to give the same band intensities as the unmodified ones.

[0218] To verify these results, the PCRs with [³²P]-labeled primers were repeated. The autoradiogram (FIG. 39) clearly shows lower yields for the modified PCR-products. The bands were excised from the gel and counted. For all PCR products the yield of the modified nucleic acids was about 50%, referring to the corresponding unmodified amplification product. Further experiments showed that exo(-)Deep-Vent and Vent DNA polymerase were able to incorporate c^7 -dATP and c^7 -dGTP during PCR as well. The overall performance, however, turned out to be best for the exo(-)Pfu DNA polymerase giving the least side products during amplification. Using all three polymerases, it was found that such PCRs employing c^7 -dATP and c^7 -dGTP instead of their isosteres showed fewer side-reactions giving a cleaner PCR-product. Decreased occurrence of amplification side products may be explained by a reduction of primer mismatches due to a lower stability of the complex formed from the primer and the 7-deaza-purine-containing template which is synthesized during PCR. Decreased melting point for DNA duplexes containing 7-deaza-purine have been described (Mizusawa, S. et al., *Nucleic Acids Res.* 14: 1319-1324

(1986)). In addition to the three polymerases specified above (exo(-) Deep Vent DNA polymerase, Vent DNA polymerase and exo(-) (Pfu) DNA polymerase), it is anticipated that other polymerases, such as the Large Klenow fragment of *E. coli* DNA polymerase, Sequenase, Taq DNA polymerase and U AmpliTaq DNA polymerase can be used. In addition, where RNA is the template, RNA polymerases, such as the SP6 or the T7 RNA polymerase, must be used.

[0219] MALDI-TOF Mass Spectrometry of Modified and Unmodified PCR Products

[0220] The 99-mer, 103-mer and 200-mer PCR products were analyzed by MALDI-TOF MS. Based on past experience, it was known that the degree of depurination depends on the laser energy used for desorption and ionization of the analyte. Since the influence of 7-deazapurine modification on fragmentation due to depurination was to be investigated, all spectra were measured at the same relative laser energy.

[0221] FIGS. 40A and 40B show the mass spectra of the modified and unmodified 103-mer nucleic acids. In case of the modified 103-mer, fragmentation causes a broad (M+H)⁺ signal. The maximum of the peak is shifted to lower masses so that the assigned mass represents a mean value of the (M+H)⁺ signal and signals of fragmented ions, rather than the (M+H)⁺ signal itself. Although the modified 103-mer still contains about 20% A and G from the oligonucleotide primers, it shows less fragmentation which is featured by much more narrow and symmetric signals. Peak tailing, especially on the lower mass side due to depurination, is substantially reduced. Hence, the difference between measured and calculated mass is strongly reduced although it is still below the expected mass. For the unmodified sample, a (M+H)⁺ signal of 31670 was observed, which is a 97 u or 0.3% difference to the calculated mass, while in the case of the modified sample, this mass difference diminished to 10 u or 0.03% (31713 u found, 31723 u calculated). These observations are verified by a significant increase in mass resolution of the (M+H)⁺ signal of the two signal strands (m/Δm=67 as opposed to 18 for the unmodified sample with Δm=full width at half maximum, fwhm). Because of the low mass difference between the two single strands (8u), their individual signals were not resolved.

[0222] With the results of the 99 base pair DNA fragments, the effects of increased mass resolution for 7-deazapurine-containing DNA becomes even more evident. The two single strands in the unmodified sample were not resolved even though the mass difference between the two strands of the PCR product was very high with 526 u due to unequal distribution of purines and pyrimidines (FIGS. 41A1 and 41A2). In contrast to this, the modified DNA showed distinct peaks for the two single strands (FIGS. 41B1 and 41B2) which makes the superiority of this approach for the determination of molecular weights to gel electrophoretic methods even more profound. Although base line resolution was not obtained, the individual masses were able to be assigned with an accuracy of 0.1% (Δm=27 u for the lighter (calc. mass=30224 u) and Δm=14 u for the heavier strand (calc. mass=30750 u)). Again, it was found that the full width at half maximum was substantially decreased for the 7-deazapurine containing sample.

[0223] In case of the 99-mer and the 103-mer, the 7-deazapurine containing nucleic acids seem to give higher sensitivity despite the fact that they still contain about 20% unmodified purine nucleotides. To get comparable signal-to-noise ratio at similar intensities for the (M+H)⁺ signals, the unmodified 99-mer required 20 laser shots in contrast to 12 for the modified one and the 103-mer required 12 shots for the unmodified sample as opposed to three for the 7-deazapurine nucleoside-containing PCR product.

[0224] Comparing the spectra of the modified and unmodified 200-mer amplicons, improved mass resolution was again found for the 7-deazapurine-containing sample as well as increased signal intensities (FIGS. 42A and 42B). While the signal of the single strands predominates in the spectrum of the modified sample the DNA-duplex and dimers of the single strands gave the strongest signal for the unmodified sample.

[0225] A complete 7-deaza purine modification of nucleic acids may be achieved either using modified primers in PCR or cleaving the unmodified primers from the partially modified PCR product. Since disadvantages are associated with modified primers, as described above, a 100-mer was synthesized using primers with a ribo-modification. The primers were cleaved hydrolytically with NaOH according to a method developed earlier in our laboratory (Köster et al., *Z. Physiol. Chem.* 359: 1579-1589 (1978)). Both hydrolyzed PCR product as well as the two released primers could be detected, together with a small signal from residual uncleaved 100-mer. This procedure is especially useful for the MALDI-TOF analysis of very short PCR-products since the share of unmodified purines originating from the primer increases with decreasing length of the amplified sequence.

[0226] The remarkable properties of 7-deazapurine modified nucleic acids can be explained by either more effective desorption and/or ionization, increased ion stability and/or a lower denaturation energy of the double stranded purine-modified nucleic acid. The exchange of the N-7 for a methine group results in the loss of one acceptor for a hydrogen bond which influences the ability of the nucleic acid to form secondary structures due to non-Watson-Crick base pairing (Seela, F. and A. Kehne, *Biochemistry* 26: 2232-2238 (1987)), which should be a reason for better desorption during the MALDI process. In addition to this, the aromatic system of 7-deazapurine has a lower electron density that weakens Watson-Crick base pairing resulting in a decreased melting point (Mizusawa, S. et al., *Nucleic Acids Res.* 14: 1319-1324 (1986)) of the double-strand. This effect may decrease the energy needed for denaturation of the duplex in the MALDI process. These aspects as well as the loss of a site which probably will carry a positive charge on the N-7 nitrogen renders the 7-deazapurine-modified nucleic acid less polar and may promote the effectiveness of desorption.

[0227] Because of the absence of N-7 as proton acceptor and the decreased polarization of the C-N bond in 7-deazapurine nucleosides, depurination following the mecha-

nisms established for hydrolysis in solution is prevented. Although a direct correlation of reactions in solution and in the gas phase is problematic, less fragmentation due to depurination of the modified nucleic acids can be expected in the MALDI process. Depurination may either be accompanied by loss of charge, which decreases the total yield of charged species, or it may produce charged fragmentation products, which decreases the intensity of the non-fragmented molecular ion signal.

[0228] The observation of both increased sensitivity and decreased peak tailing of the (M+H)⁺ signals on the lower mass side due to decreased fragmentation of the 7-deazapurine-containing samples indicate that the N-7 atom indeed is essential for the mechanism of depurination in the MALDI-TOF process. In conclusion, 7-deazapurine-containing nucleic acids show distinctly increased ion-stability and sensitivity under MALDI-TOF conditions and therefore provide for higher mass accuracy and mass resolution.

Sequencing complex:
5'-AAGATCTGACCAGGATTTCGGTTAGCGTGA CTGCTGCTGCTGCTGCTGC (SEQ.ID.No.25)
TGGATGATCCGACGCATCAGATCTGG-(A²)_n-3'
(TNR.PLASM2)
3'-CTACTAGGCTGCGTAGTC-5' (CM1) (SEQ.ID.No.26)

EXAMPLE 9

Solid State Sequencing and Mass Spectrometer Detection

Materials and Methods

[0229] Oligonucleotides were purchased from Operon Technologies (Alameda, Calif.) in an unpurified form. Sequencing reactions were performed on a solid surface using reagents from the sequencing kit for Sequenase Version 2.0 (Amersham, Arlington Heights, Ill.).

Sequencing a 39-mer target
Sequencing complex:
5'-TCTGGCCCTGGTGCAGGGCCTATTGTAGTTGTGACGTACA-(A²)_n-3' (SEQ.ID.No.23)
(DNA 11683)
3'-TCAACACTGCATGT-5' (PNA 16/DNA) (SEQ.ID.No.24)

[0230] In order to perform solid-state DNA sequencing, template strand DNA11683 was 3'-biotinylated by terminal deoxynucleotidyl transferase. A 30 μ l reaction, containing 60 pmol of DNA11683, 1.3 nmol of biotin 14-dATP (GIBCO BRL, Grand Island, N.Y.), 30 units of terminal transferase (Amersham, Arlington Heights, Ill.), and 1x reaction buffer (supplied with enzyme), was incubated at 37° C. for 1 hour. The reaction was stopped by heat inactivation of the terminal transferase at 70° C. for 10 min. The resulting product was desalted by passing through a TE-10 spin column (Clontech). More than one molecule of biotin-14-dATP could be added to the 3'-end of DNA11683. The biotinylated DNA11683 was incubated with 0.3 mg of Dynal streptavidin beads in 30 μ l 1x binding and washing buffer at ambient temperature for 30 min. The beads were washed twice with TE, resuspended in 30 μ l TE, and a 10 μ l aliquot (containing 0.1 mg of beads) was used for sequencing reactions.

[0231] The 0.1 mg beads from previous step were resuspended in a 10 μ l volume containing 2 μ l of 5x Sequenase buffer (200 mM Tris-HCl, pH 7.5, 100 mM MgCl₂, and 250 mM NaCl) from the Sequenase kit and 5 pmol of corresponding primer PNA16/DNA. The annealing mixture was heated to 70° C. and allowed to cool slowly to room temperature over a 20-30 min. time period. Then 1 μ l 0.1 M dithiothreitol solution, 1 μ l Mn buffer (0.15 M sodium isocitrate and 0.1 M MnCl₂), and 2 μ l of diluted Sequenase (3.25 units) were added. The reaction mixture was divided into four aliquots of 3 μ l each and mixed with termination mixes (each consists of 3 μ l of the appropriate termination mix: 32 μ M c7dATP, 32 μ M dCTP, 32 μ M c7dGTP, 32 μ M dTTP and 3.2 μ M of one of the four ddTNPS, in 50 mM NaCl). The reaction mixtures were incubated at 37° C. for 2 min. After the completion of extension, the beads were precipitated and the supernatant was removed. The beads were washed twice and resuspended in TE and kept at 4° C.

[0232] Sequencing a 78-mer Target

[0233] The target TNR.PLASM2 was biotinylated and sequenced using procedures similar to those described in previous section (sequencing a 39-mer target).

[0234] Sequencing a 15-mer Target With Partially-Duplex Probe

Sequencing complex:

-continued (SEQ.ID.No.27)
5'-F-GATGATCCGACGCATCAGCTC-3'
(SEQ.ID.No.28)
3'-b-CTACTAGGCTGCGTAGTCTCGAGAACCCTGGCT-5'

[0235] CM1B3B was immobilized on DynaBeads M280 with streptavidin (Dynal, Norway) by incubating 60 pmol of CM1B3B with 0.3 mg magnetic beads in 30 μ l 1 M NaCl and TE (1x binding and washing buffer) at room temperature for 30 min. The beads were washed twice with TE, resuspended in 30 μ l TE, and a 10 or 20 μ l aliquot (containing 0.1 or 0.2 mg of beads respectively) was used for sequencing reactions.

[0236] The duplex was formed by annealing a corresponding aliquot of beads from previous step with 10 pmol of

DF11a5F (or 20 pmol of DF11a5F for 0.2 mg of beads) in a 9 μ l volume containing 2 μ l of 5 \times Sequenase buffer (200 mM Tris-HCl, pH 7.5, 100 mM MgCl₂, and 250 mM NaCl) from the Sequenase kit. The annealing mixture was heated to 65° C. and allowed to cool slowly to 37° C. over a 20-30 min time period. The duplex primer was then mixed with 10 pmol of TS10 (20 pmol of TS10 for 0.2 mg of beads) in 1 μ l volume, and the resulting mixture was further incubated at 37° C. for 5 min., and room temperature for 5-10 min. Then 1 μ l 0.1 M dithiothreitol solution, 1 μ l Mn buffer (0.15 M sodium isocitrate and 0.1 M MnCl₂), and 2 μ l of diluted Sequenase (3.25 units) were added. The reaction mixture was divided into four aliquots of 3 μ l each and mixed with termination mixes (each consists of 4 μ l of the appropriate termination mix: 16 μ M dATP, 16 μ M dCTP, 16 μ M dGTP, 16 μ M dTTP and 1.6 μ M of one of the four ddNTPs, in 50 mM NaCl). The reaction mixtures were incubated at room temperature for 5 min., and 37° C. for 5 min. After the completion of extension, the beads were precipitated and the supernatant was removed. The beads were resuspended in 20 μ l TE and kept at 4° C. An aliquot of 2 μ l (out of 20 μ l) from each tube was taken and mixed with 8 μ l of formamide, the resulting samples were denatured at 90-95° C. for 5 min. and 2 μ l (out of 10 μ l total) was applied to an ALF DNA sequencer (Pharmacia, Piscataway, N.J.) using a 10% polyacrylamide gel containing 7 M urea and 0.6 \times TBE. The remaining aliquot was used for MALDI-TOF MS analysis.

[0237] MALDI Sample Preparation and Instrumentation

[0238] Before MALDI analysis, the sequencing-ladder-loaded magnetic beads were washed twice using 50 mM ammonium citrate and resuspended in 0.5 μ l pure water. The suspension was then loaded onto the sample target of the mass spectrometer and 0.5 μ l of saturated matrix solution (3-hydroxyisobutyric acid (HPA):ammonium citrate=10:1 mole ratio in 50% acetonitrile) was added. The mixture was allowed to dry prior to mass spectrometer analysis.

[0239] The reflectron TOFMS mass spectrometer (Vision 2000, Finnigan MAT, Bremen, Germany) was used for analysis. 5 kV was applied in the ion source and 20 kV was applied for postacceleration. All spectra were taken in the positive ion mode and a nitrogen laser was used. Normally, each spectrum was averaged for more than 100 shots and a standard 25-point smoothing was applied.

Results and Discussion

[0240] Conventional Solid-State Sequencing

[0241] In conventional sequencing methods, a primer is directly annealed to the template and then extended and terminated in a Sanger dideoxy sequencing. Normally, a biotinylated primer is used and the sequencing ladders are captured by streptavidin-coated magnetic beads. After washing, the products are eluted from the beads using EDTA and formamide. However, our previous findings indicated that only the annealed strand of a duplex is desorbed and the immobilized strand remains on the beads. Therefore, it is advantageous to immobilize the template and anneal the primer. After the sequencing reaction and washing, the beads with the immobilized template and annealed sequencing ladder can be loaded directly onto the mass spectrometer target and mixed with matrix. In MALDI, only the annealed sequencing ladder will be desorbed and ionized, and the immobilized template will remain on the target.

[0242] A 39-mer template (SEQ.ID.No. 23) was first biotinylated at the 3' end by adding biotin-14-dATP with terminal transferase. More than one biotin-14-dATP molecule could be added by the enzyme. However, since the template was immobilized and remained on the beads during MALDI, the number of biotin-14-dATP added would not affect the mass spectra. A 14-mer primer (SEQ.ID.No. 29) was used for the solid-state sequencing. MALDI-TOF mass spectra of the four sequencing ladders are shown in FIGS. 44A-44D and the expected theoretical values are shown in Table II.

TABLE II

1	5'-TCTGGCCTGGTGCAGGGCCTATTGTAGTGTGACGTACA-(A ^b) ₂₅ -3'
2	3'-TCACACTGCATGT-5'
3	3'-ATCAACACTGCATGT-5'
4	3'-CATCAACACTGCATGT-5'
5	3'-ACATCAACACTGCATGT-5'
6	3'-AACATCAACACTGCATGT-5'
7	3'-TAACATCAACACTGCATGT-5'
8	3'-ATAACATCAACACTGCATGT-5'
9	3'-GATAACATCAACACTGCATGT-5'
10	3'-GGATAACATCAACACTGCATGT-5'
11	3'-CGGATAACATCAACACTGCATGT-5'
12	3'-CCGGATAACATCAACACTGCATGT-5'
13	3'-CCCCGATAACATCAACACTGCATGT-5'
14	3'-TCCCCGATAACATCAACACTGCATGT-5'
15	3'-GTCCCCGATAACATCAACACTGCATGT-5'

TABLE II-continued

16	3'-CGTCCCGGATAACATCAACACTGCATGT-5'			
17	3'-ACGTCCCGGATAACATCAACACTGCATGT-5'			
18	3'-CAGTCCCGGATAACATCAACACTGCATGT-5'			
19	3'-CCACGTCCCGGATAACATCAACACTGCATGT-5'			
20	3'-ACCACGTCCCGGATAACATCAACACTGCATGT-5'			
21	3'-GACCACGTCCCGGATAACATCAACACTGCATGT-5'			
22	3'-GGACCACGTCCCGGATAACATCAACACTGCATGT-5'			
23	3'-CGGACCACGTCCCGGATAACATCAACACTGCATGT-5'			
24	3'-CCGGACCACGTCCCGGATAACATCAACACTGCATGT-5'			
25	3'-ACCGGACCACGTCCCGGATAACATCAACACTGCATGT-5'			
26	3'-GACCGGACCACGTCCCGGATAACATCAACACTGCATGT-5'			
27	3'-AGACCGGACCACGTCCCGGATAACATCAACACTGCATGT-5'			
	A-reaction	C-reaction	G-reaction	T-reaction
1.				
2.	4223.8	4223.8	4223.8	4223.8
3.	4521.1			
4.		4810.2		
5.	5123.4			
6.	5436.6			
7.				
8.	6054.0			5740.8
9.				
10.			6383.2	
11.		7001.6	6712.4	
12.		7290.8		
13.		7580.0		
14.				7884.2
15.			8213.4	
16.		8502.6		
17.	8815.8			
18.		9105.0		
19.		9394.2		
20.	9707.4			
21.			10036.6	
22.			10365.8	
23.		10655.0		
24.		10944.2		
25.	11257.4			
26.			11586.6	
27.	11899.8			

[0243] The sequencing reaction produced a relatively homogenous ladder, and the full-length sequence was determined easily. One peak around 5150 appeared in all reactions was not identified. A possible explanation is that a small portion of the template formed some kind of secondary structure, such as a loop, which hindered Sequenase extension. Mis-incorporation is of minor importance, since the intensity of these peaks were much lower than that of the sequencing ladders. Although 7-deaza purines were used in the sequencing reaction, which could stabilize the N-glycosidic bond and prevent depurination, minor base losses were still observed since the primer was not substituted by 7-deazapurines. The full length ladder, with a ddA at the 3' end, appeared in the A reaction with an apparent mass of 11899.8. However, a more intense peak of 122 appeared in

all four reactions and is likely due to an addition of an extra nucleotide by the Sequenase enzyme.

[0244] The same technique could be used to sequence longer DNA fragments. A 78-mer template containing a CTG repeat (SEQ.ID.No. 25) was 3'-biotinylated by adding biotin-14-dATP with terminal transferase. An 18-mer primer (SEQ.ID.No. 26) was annealed right outside the CTG repeat so that the repeat could be sequenced immediately after primer extension. The four reactions were washed and analyzed by MALDI-TOF MS as usual. An example of the G-reaction is shown in FIG. 45 and the expected sequencing ladder is shown in Table III with theoretical mass values for each ladder component. All sequencing peaks were well resolved except the last component (theoretical value 20577.4) was indistinguishable from the background. Two

neighboring sequencing peaks (a 62-mer and a 63-mer) were also separated indicating that such sequencing analysis could be applicable to longer templates. Again, an addition of an extra nucleotide by the Sequenase enzyme was observed in this spectrum. This addition is not template

specific and appeared in all four reactions which makes it easy to be identified. Compared to the primer peak, the sequencing peaks were at much lower intensity in the long template case. Further optimization of the sequencing reaction may be required.

TABLE III

AAGATCTGACCAGGGATTCGGTTAGCGTGA CTGCTGCTGCTGGATGATCCGACGCATCAGATCTGG- (A ⁿ) _n -3'	
1	3'-CTACTAGGCTGCGTAGTC-5'
2	3'-CCTACTAGGCTGCGTAGTC-5'
3	3'-ACCTACTAGGCTGCGTAGTC-5'
4	3'-GACCTACTAGGCTGCGTAGTC-5'
5	3'-CGACCTACTAGGCTGCGTAGTC-5'
6	3'-ACGACCTACTAGGCTGCGTAGTC-5'
7	3'-GACGACCTACTAGGCTGCGTAGTC-5'
8	3'-CGACGACCTACTAGGCTGCGTAGTC-5'
9	3'-ACGACGACCTACTAGGCTGCGTAGTC-5'
10	3'-GACGACGACCTACTAGGCTGCGTAGTC-5'
11	3'-CGACGACGACCTACTAGGCTGCGTAGTC-5'
12	3'-ACGACGACGACCTACTAGGCTGCGTAGTC-5'
13	3'-GACGACGACGACCTACTAGGCTGCGTAGTC-5'
14	3'-CGACGACGACGACCTACTAGGCTGCGTAGTC-5'
15	3'-ACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
16	3'-GACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
17	3'-CGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
18	3'-ACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
19	3'-GACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
20	3'-CGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
21	3'-ACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
22	3'-GACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
23	3'-CGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
24	3'-ACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
25	3'-GACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
26	3'-TCGACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
27	3'-CTGACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
28	3'-ACTGACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
29	3'-CACTGACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
30	3'-GCACTGACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
31	3'-CGCACTGACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
32	3'-TCGCACTGACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'

TABLE III—continued

33	3'-ATCGCACTGACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
34	3'-AATCGCACTGACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
35	3'-CAATCGCACTGACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
36	3'-CCAATCGCACTGACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
37	3'-GCCAATCGCACTGACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
38	3'-AGCCAATCGCACTGACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
39	3'-AAGCCAATCGCACTGACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
40	3'-TAAGCCAATCGCACTGACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
41	3'-CTAAGCCAATCGCACTGACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
42	3'-CCTAAGCCAATCGCACTGACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
43	3'-CCCTAAGCCAATCGCACTGACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
44	3'-TCCCTAAGCCAATCGCACTGACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
45	3'-GTCCCTAAGCCAATCGCACTGACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
46	3'-GGTCCCTAAGCCAATCGCACTGACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
47	3'-TGGTCCCTAAGCCAATCGCACTGACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
48	3'-CTGGTCCCTAAGCCAATCGCACTGACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
49	3'-ACTGGTCCCTAAGCCAATCGCACTGACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
50	3'-GACTGGTCCCTAAGCCAATCGCACTGACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
51	3'-AGACTGGTCCCTAAGCCAATCGCACTGACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
52	3'-TAGACTGGTCCCTAAGCCAATCGCACTGACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
53	3'-CTAGACTGGTCCCTAAGCCAATCGCACTGACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
54	3'-TCTAGACTGGTCCCTAAGCCAATCGCACTGACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'
55	3'-TTCTAGACTGGTCCCTAAGCCAATCGCACTGACGACGACGACGACGACGACGACCTACTAGGCTGCGTAGTC-5'

	ddATP	ddCTP	ddGTP	ddTTP
1.	5491.6	5491.6	5491.6	5491.6
2.		5764.8		
3.	6078.0			
4.			6407.2	
5.		6696.4		
6.	7009.6			
7.			7338.8	
8.		7628.0		
9.	7941.2			
10.			8270.4	
11.		8559.6		
12.	8872.8			
13.			9202.0	
14.		9491.2		
15.	9804.4			
16.			10133.6	
17.		10422.88		
18.	10736.0			
19.		11354.4	11065.2	
20.				11996.8
21.	11667.6			
22.		12286.0		
23.				
24.	12599.2			
25.			12928.4	
26.				13232.6
27.		13521.8		

TABLE III—continued

28.	13835.0			
29.		14124.2		
30.			14453.4	
31.		14742.6		
32.				15046.8
33.	15360.0			
34.	15673.2			
35.		15962.4		
36.		16251.6		
37.			16580.8	
38.	16894.0			
39.	17207.2			
40.				17511.4
41.		17800.6		
42.		18189.8		
43.		18379.0		
44.				18683.2
45.			19012.4	
46.			19341.6	
47.				19645.8
48.		19935.0		
49.	20248.2			
50.			20577.4	
51.	20890.6			
52.				21194.4
53.		21484.0		
54.				21788.2
55.				22092.4

[0245] Sequencing Using Duplex DNA Probes for Capturing and Priming

[0246] Duplex DNA probes with single-stranded overhang have been demonstrated to be able to capture specific DNA templates and also served as primers for solid-state sequencing. The scheme is shown in FIG. 46. Stacking interactions between a duplex probe and a single-stranded template allow only 5-base overhang to be to be sufficient for capturing. Based on this format, a 5' fluorescent-labeled 23-mer (5'-GAT GAT CCG ACG CAT CAC AGC TC-3') (SEQ.ID.No. 29)) was annealed to a 3'-biotinylated 18-mer (5'-GTG ATG CCT CGG ATC ATC-3') (SEQ.ID.No. 30)), leaving a 5-base overhang. A 15-mer template (5'-TCG GTT CCA AGA GCT-3') (SEQ.ID.No. 31)) was captured by the duplex and sequencing reactions were performed by extension of the 5-base overhang. MALDI-TOF mass spectra of the reactions are shown in FIGS. 47A-47D. All sequencing peaks were resolved, although at relatively low intensities. The last peak in each reaction is due to unspecific addition of one nucleotide to the full length extension product by the Sequenase enzyme. For comparison, the same products were run on a conventional DNA sequencer and a stacking fluorogram of the results is shown in FIGS. 48A-48D. As can be seen from the Figures, the mass spectra had the same pattern as the fluorogram with sequencing peaks at much lower intensity compared to the 23-mer primer.

[0247] Improvements of MALDI-TOF Mass Spectrometry as a Detection Technique

[0248] Sample distribution can be made more homogeneous and signal intensity could potentially be increased by implementing the picoliter vial technique. In practice, the samples can be loaded on small pits with square openings of 100 μm size. The beads used in the solid state sequencing are

less than 10 μm in diameter, so they should fit well in the microliter vials. Microcrystals of matrix and DNA containing "sweet spots" will be confined in the vial. Since the laser spot size is about 100 μm in diameter, it will cover the entire opening of the vial. Therefore, searching for sweet spots will be unnecessary and high repetition-rate laser (e.g. ≤ 10 Hz) can be used for acquiring spectra. An earlier report has shown that this device is capable of increasing the detection sensitivity of peptides and proteins by several orders of magnitude compared to the conventional MALDI sample preparation technique.

[0249] Resolution of MALDI on DNA needs to be further improved in order to extend the sequencing range beyond 100 bases. Currently, using 3-HPA/ammonium citrate as matrix and a reflectron TOF mass spectrometer with 5 kV ion source and 20 kV postacceleration, the resolution of the run-through peak in FIG. 33A (73-mer) is greater than 200 (FWHM) which is enough for sequence determination in this case. This resolution is also the highest reported for MALDI desorbed DNA ions above the 70-mer range. Use of the delayed extraction technique may further enhance resolution.

[0250] All of the above-cited references, applications and publications are herein incorporated by reference.

Equivalents

[0251] Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, numerous equivalents to the specific procedures described herein. Such equivalents are considered to be within the scope of this invention and are covered by the following claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(iii) NUMBER OF SEQUENCES: 33

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

GCAAGTGAAT CCTGAGCGTG

20

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 19 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

GTGTGAAGGG TTCATATGC

19

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 28 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

ATCTATATTC ATCATAGGAA ACACCACA

28

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 30 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

GTATCTATAT TCATCATAGG AAACACCAAT

30

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 30 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

-continued

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

GCTTTGGGGC ATGCACATTC ACCCGTATAA 30

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 30 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

CTGACTACTA ATTCCTTGA TGCTGGGTCT 30

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

TTGCCTGAGT GCACATATGCT 20

(2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

AGCTCTATAT CGGAAGCCT 20

(2) INFORMATION FOR SEQ ID NO: 9:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

TTGTGCCACG CGGTGGGAA TGTA 24

(2) INFORMATION FOR SEQ ID NO: 10:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 25 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

-continued

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:
AGCAACGACT GTTGCCCGC CAGTTG 26

(2) INFORMATION FOR SEQ ID NO: 11:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 25 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:
TACATCCCA ACCCGGTGC ACAAC 25

(2) INFORMATION FOR SEQ ID NO: 12:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 25 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:
AACTGGCGGG CAAACAGTCG TTGCT 25

(2) INFORMATION FOR SEQ ID NO: 13:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 57 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:
ACCATTAAG AAAATATCAT CTTTGGTGTT TCCTATGATG AATATAGAAG CGTCATC 57

(2) INFORMATION FOR SEQ ID NO: 14:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 24 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:
ACCACAAAGG ATACTACTTA TATC 24

(2) INFORMATION FOR SEQ ID NO: 15:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 29 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

-continued

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:
TAGAAACCAC AAAGGATACT ACTTATATC 29

(2) INFORMATION FOR SEQ ID NO: 16:
(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 26 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
(ii) MOLECULE TYPE: cDNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:
TAACCACAAA GGATACTACT TATATC 26

(2) INFORMATION FOR SEQ ID NO: 17:
(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 29 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
(ii) MOLECULE TYPE: cDNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:
TAGAAACCAC AAAGGATACT ACTTATATC 29

(2) INFORMATION FOR SEQ ID NO: 18:
(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 38 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
(ii) MOLECULE TYPE: cDNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:
CTTTTATAGT AGAAACCACA AAGGATACTA CTTATATC 38

(2) INFORMATION FOR SEQ ID NO: 19:
(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 35 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
(ii) MOLECULE TYPE: cDNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:
CTTTTATAGT AACCACAAG GATACTACTT ATATC 35

(2) INFORMATION FOR SEQ ID NO: 20:
(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 35 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
(ii) MOLECULE TYPE: cDNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:

-continued

CTTTTATAGA AACCACAAAG GATACTACTT ATATC 35

(2) INFORMATION FOR SEQ ID NO: 21:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 31 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 21:

CGTAGAAACC ACAAGGATA CTAATTATAT C 31

(2) INFORMATION FOR SEQ ID NO: 22:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 58 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22:

GAATTACATT CCCAACC GCG TGGCACAACA ACTGGCGGGC AAACAGTCGT TGCTGATT 58

(2) INFORMATION FOR SEQ ID NO: 23:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 58 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 23:

AATCAGCAAC GACTGTTTGC CCGCCAGTTG TTGTGCCACG CGGTGGGAA TGTAAATTC 58

(2) INFORMATION FOR SEQ ID NO: 24:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 252 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 24:

GGCACGGCTG TCCAAGGAGC TGCAGGCGGC GCAGGCCCGG CTGGCGCGCG ACATGGAGGA 60

CGTGTGCGCC GCTGTGTGA GTACCGCGGC GAGGTGCAGG CCATGCTCGG CCAGAGCAC 120

GAGGAGCTGC GGTGCGCCT CGCCTCCAC CTGCGCAAGC TGCCTAAGCG GCTCCTCCG 180

GATGCGGATG ACCTGCAGAA GTCCCTGGCA GTGTACCAGG CCGGGGCCCG CGAGGGCGC 240

GAGCGCGGCC TC 252

(2) INFORMATION FOR SEQ ID NO: 25:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 110 base pairs

-continued

(B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 25:

GCAACATTTT GCTCCCGGTC ACGGTTCCAA CGTACGGACG TCCAGCTGAG ATCTCCTAGG 60
 GGCCCATGGC TCGAGCTTAA GCATTAGTAC CAGTATCGAC AAAGGACACA 110

(2) INFORMATION FOR SEQ ID NO: 26:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 110 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 26:

TGTGTCCTTT GTCGATACTG GTACTAATGC TTAAGCTCGA GCCATGGGCC CCTAGGAGAT 60
 CTCAGCTGGA CGTCCGTACG TTCCAACCGT GACCGGCAGC AAAATGTTGC 110

(2) INFORMATION FOR SEQ ID NO: 27:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 217 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 27:

AACGTGCTGC CTCCACCGC GATGTTGATG ATTATGTGTC TGAATTGAT GGGGGCAGGC 60
 GGCCCCCGTC TGTTTGTGCG GGGTCTGGTG TTGATGGTGG TTTCCTGCCT TGTACACCT 120
 GACCTGCAGC CCAAGCTTGG GATCCACCAC CATCACCATC ACTAATAATG CATGGGGCTG 180
 AGCCAATFTGG CACTGGCCGT CGTTTACAA CGTCGTG 217

(2) INFORMATION FOR SEQ ID NO: 28:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 217 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 28:

CACGACGTTG TAAACGACG GCCAGTGCCA ATTGGCTGCA GCCCATGCAT TATTAGTGAT 60
 GGTGATGGTG GTGATCCCA AGCTTGGGCT GCAGGTCGAG GGTGACAAGG CAGGAAGC 120
 CCATCAACAC CAGACCCGGG ACAACAGAC GGGGGCCGCC TGCCCCCATC AAATTCAGA 180
 ACATAATCAT CAACATCGCG GTGGAAGGCA GCACGTT 217

(2) INFORMATION FOR SEQ ID NO: 29:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 17 base pairs

-continued

(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 29:

GTAAAACGAC GGCCAGT 17

(2) INFORMATION FOR SEQ ID NO: 30:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 17 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 30:

CAGGAAACAG CTATGAC 17

(2) INFORMATION FOR SEQ ID NO: 31:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 31:

CTTCCACCGC GATGTTGA 18

(2) INFORMATION FOR SEQ ID NO: 32:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 19 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 32:

TTGTAAAACG ACGCCCACT 19

(2) INFORMATION FOR SEQ ID NO: 33:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 33:

GTCACCCTCG ACCTGCAG 18

What is claimed is:

1. A method for detecting mutations in a target nucleic acid, comprising:

digesting a target nucleic acid molecule;

capturing digested fragments on a solid support that comprises oligonucleotides complementary thereto;

detecting hybrids and the molecular weights of captured fragments by mass spectrometry, thereby identifying mutations in the target nucleic acid molecule.

2. The method of claim 1, wherein the target molecule is RNA.

3. The method of claim 2, wherein the RNA is produced by transcription of a target DNA molecule.

4. The method of claim 1, wherein the solid support comprises a flat surface.

5. The method of claim 1, wherein the solid support comprises matrix for performing matrix-assisted laser desorption/ionization (MALDI) mass spectrometry.

6. The method of claim 1, wherein the solid support is selected from the group consisting of glass fiber filters, glass surfaces and metal surfaces.

7. The method of claim 1, wherein the solid support is selected from the group consisting of steel, gold, silver, aluminum, copper and silicon.

8. The method of claim 1, wherein the solid support is silicon.

9. The method of claim 1, wherein the solid support is a silicon wafer.

10. The method of claim 1, wherein the oligonucleotides on the solid support are linked to the support via a linker or a bond cleavable under the conditions of mass spectrometric analysis,

11. The method of claim 10, wherein the oligonucleotides are linked to the support via a linker or bond is photocleavable.

12. The method of claim 1, wherein the oligonucleotide comprises oligoribonucleotides, oligodeoxyribonucleotides, nucleotide analogs, or protein nucleic acid (PNA).

13. The method of claim 10, wherein the nucleotide analogs comprise a thio-modified phosphodiester or phosphotriester backbone.

14. The method of claim 1, wherein the oligonucleotides are linked to the support by an irreversible bond.

15. The method of claim 1, wherein the oligonucleotides are linked to the support by a disulfide bond, a biotin/streptavidin linkage, a heterobifunctional derivative of a trityl ether group, a leuvinyl group, an arginine-arginine bond, a lysine-lysine bond, a pyrophosphate bond and a charge transfer complex.

* * * * *